

表 1299 Non-HDL-C濃度(mg/dL) 医療機関別 (施設血液透析患者 透析回数週3回)

医療機関	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
国公立大学	7	18	31	40	24	9	5	6	3		143	289	432	100.25	34.71
(%)	(4.9)	(12.6)	(21.7)	(28.0)	(16.8)	(6.3)	(3.5)	(4.2)	(2.1)		(100.0)				
私立大学	23	82	160	189	148	93	50	15	11	13	784	773	1,557	107.05	36.62
(%)	(2.9)	(10.5)	(20.4)	(24.1)	(18.9)	(11.9)	(6.4)	(1.9)	(1.4)	(1.7)	(100.0)				
国立病院機構等	8	11	41	39	29	25	11	7	5	2	178	278	456	109.28	39.86
(%)	(4.5)	(6.2)	(23.0)	(21.9)	(16.3)	(14.0)	(6.2)	(3.9)	(2.8)	(1.1)	(100.0)				
県市町村立・国保	187	897	1,697	1,819	1,387	718	386	156	66	58	7,371	5,152	12,523	102.95	33.87
(%)	(2.5)	(12.2)	(23.0)	(24.7)	(18.8)	(9.7)	(5.2)	(2.1)	(0.9)	(0.8)	(100.0)				
JCHO(旧社会保険)等	27	171	339	368	317	216	97	56	18	17	1,626	821	2,447	108.03	35.62
(%)	(1.7)	(10.5)	(20.8)	(22.6)	(19.5)	(13.3)	(6.0)	(3.4)	(1.1)	(1.0)	(100.0)				
厚生連等	85	414	957	1,052	872	585	253	129	58	37	4,442	1,960	6,402	107.58	34.13
(%)	(1.9)	(9.3)	(21.5)	(23.7)	(19.6)	(13.2)	(5.7)	(2.9)	(1.3)	(0.8)	(100.0)				
その他公的	80	408	897	1,027	802	467	235	111	33	18	4,078	2,465	6,543	105.28	32.56
(%)	(2.0)	(10.0)	(22.0)	(25.2)	(19.7)	(11.5)	(5.8)	(2.7)	(0.8)	(0.4)	(100.0)				
私立総合	75	358	775	869	664	397	209	88	24	23	3,482	1,376	4,858	105.13	33.37
(%)	(2.2)	(10.3)	(22.3)	(25.0)	(19.1)	(11.4)	(6.0)	(2.5)	(0.7)	(0.7)	(100.0)				
私立	970	4,326	8,628	9,926	7,720	4,745	2,395	1,063	436	260	40,469	21,615	62,084	105.58	33.99
(%)	(2.4)	(10.7)	(21.3)	(24.5)	(19.1)	(11.7)	(5.9)	(2.6)	(1.1)	(0.6)	(100.0)				
私立診療所	1,083	6,382	15,218	18,661	15,238	9,381	4,892	2,098	789	540	74,282	37,952	112,234	108.49	33.25
(%)	(1.5)	(8.6)	(20.5)	(25.1)	(20.5)	(12.6)	(6.6)	(2.8)	(1.1)	(0.7)	(100.0)				
総計	2,545	13,067	28,743	33,990	27,201	16,636	8,533	3,729	1,443	968	136,855	72,681	209,536	107.10	33.62
(%)	(1.9)	(9.5)	(21.0)	(24.8)	(19.9)	(12.2)	(6.2)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4624##

(C)Japanese Society for Dialysis Therapy

2016年調査