

表 1303 Non-HDL-C濃度(mg/dL) 原疾患別(血液透析濾過患者 透析回数週3回)

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	226	1,245	2,991	4,008	3,452	2,140	1,121	441	191	90	15,905	6,483	22,388	109.89	32.77
(%)	(1.4)	(7.8)	(18.8)	(25.2)	(21.7)	(13.5)	(7.0)	(2.8)	(1.2)	(0.6)	(100.0)				
慢性腎盂腎炎	5	25	86	140	83	53	31	18	4	3	448	179	627	109.77	31.90
(%)	(1.1)	(5.6)	(19.2)	(31.3)	(18.5)	(11.8)	(6.9)	(4.0)	(0.9)	(0.7)	(100.0)				
急速進行性糸球体腎炎	4	24	46	95	69	58	25	14	5	7	347	137	484	115.85	36.94
(%)	(1.2)	(6.9)	(13.3)	(27.4)	(19.9)	(16.7)	(7.2)	(4.0)	(1.4)	(2.0)	(100.0)				
妊娠腎/妊娠中毒症	2	12	55	80	66	56	41	12	2	3	329	130	459	117.94	32.58
(%)	(0.6)	(3.6)	(16.7)	(24.3)	(20.1)	(17.0)	(12.5)	(3.6)	(0.6)	(0.9)	(100.0)				
その他の分類不能の腎炎	2	16	51	57	52	35	15	5	2	1	236	119	355	108.85	30.47
(%)	(0.8)	(6.8)	(21.6)	(24.2)	(22.0)	(14.8)	(6.4)	(2.1)	(0.8)	(0.4)	(100.0)				
多発性嚢胞腎	17	138	361	458	422	245	114	36	12	8	1,811	758	2,569	108.70	30.83
(%)	(0.9)	(7.6)	(19.9)	(25.3)	(23.3)	(13.5)	(6.3)	(2.0)	(0.7)	(0.4)	(100.0)				
腎硬化症	68	314	865	1,050	858	540	225	108	45	20	4,093	1,803	5,896	107.68	32.15
(%)	(1.7)	(7.7)	(21.1)	(25.7)	(21.0)	(13.2)	(5.5)	(2.6)	(1.1)	(0.5)	(100.0)				
悪性高血圧	4	24	67	94	85	48	30	16	8	3	379	176	555	113.27	34.45
(%)	(1.1)	(6.3)	(17.7)	(24.8)	(22.4)	(12.7)	(7.9)	(4.2)	(2.1)	(0.8)	(100.0)				
糖尿病性腎症	348	1,771	3,803	4,064	3,244	1,935	1,016	422	182	102	16,887	7,629	24,516	105.47	33.51
(%)	(2.1)	(10.5)	(22.5)	(24.1)	(19.2)	(11.5)	(6.0)	(2.5)	(1.1)	(0.6)	(100.0)				
SLE腎炎	8	31	67	85	88	42	38	13	2	1	375	162	537	110.46	33.40
(%)	(2.1)	(8.3)	(17.9)	(22.7)	(23.5)	(11.2)	(10.1)	(3.5)	(0.5)	(0.3)	(100.0)				
アミロイド腎	1	4	13	17	9	7	6	5			62	36	98	110.71	34.91
(%)	(1.6)	(6.5)	(21.0)	(27.4)	(14.5)	(11.3)	(9.7)	(8.1)			(100.0)				
痛風腎	2	21	32	49	30	19	10	4	2	1	170	57	227	105.97	33.65
(%)	(1.2)	(12.4)	(18.8)	(28.8)	(17.6)	(11.2)	(5.9)	(2.4)	(1.2)	(0.6)	(100.0)				
先天性代謝異常による腎不全	2	1	10	9	12	7	2	2		1	46	27	73	111.50	37.31
(%)	(4.3)	(2.2)	(21.7)	(19.6)	(26.1)	(15.2)	(4.3)	(4.3)		(2.2)	(100.0)				
腎・尿路結核	1	1	6	7	9		2	1			27	17	44	106.37	29.32
(%)	(3.7)	(3.7)	(22.2)	(25.9)	(33.3)		(7.4)	(3.7)			(100.0)				
腎・尿路結石	1	9	12	17	20	8	3	3	1		74	39	113	107.09	31.72
(%)	(1.4)	(12.2)	(16.2)	(23.0)	(27.0)	(10.8)	(4.1)	(4.1)	(1.4)		(100.0)				
腎・尿路腫瘍	3	18	21	39	26	19	9	1	1		137	53	190	104.58	30.64
(%)	(2.2)	(13.1)	(15.3)	(28.5)	(19.0)	(13.9)	(6.6)	(0.7)	(0.7)		(100.0)				
閉塞性尿路障害		5	16	30	33	15	10	2	1		112	39	151	114.21	27.57
(%)		(4.5)	(14.3)	(26.8)	(29.5)	(13.4)	(8.9)	(1.8)	(0.9)		(100.0)				
骨髄腫	1	5	4	6	3	1	3	2			25	12	37	103.08	41.18
(%)	(4.0)	(20.0)	(16.0)	(24.0)	(12.0)	(4.0)	(12.0)	(8.0)			(100.0)				
腎形成不全	3	10	17	33	36	8	9	6	1	3	126	45	171	112.96	35.97
(%)	(2.4)	(7.9)	(13.5)	(26.2)	(28.6)	(6.3)	(7.1)	(4.8)	(0.8)	(2.4)	(100.0)				
不明	95	368	865	996	899	594	286	127	50	24	4,304	2,227	6,531	108.59	33.93
(%)	(2.2)	(8.6)	(20.1)	(23.1)	(20.9)	(13.8)	(6.6)	(3.0)	(1.2)	(0.6)	(100.0)				
再導入	7	33	87	108	86	63	38	14	8	4	448	185	633	112.12	35.48
(%)	(1.6)	(7.4)	(19.4)	(24.1)	(19.2)	(14.1)	(8.5)	(3.1)	(1.8)	(0.9)	(100.0)				
その他	20	90	242	285	231	152	90	25	7	7	1,149	571	1,720	108.15	32.63
(%)	(1.7)	(7.8)	(21.1)	(24.8)	(20.1)	(13.2)	(7.8)	(2.2)	(0.6)	(0.6)	(100.0)				
合計	820	4,165	9,717	11,727	9,813	6,045	3,124	1,277	524	278	47,490	20,884	68,374	108.05	33.15
(%)	(1.7)	(8.8)	(20.5)	(24.7)	(20.7)	(12.7)	(6.6)	(2.7)	(1.1)	(0.6)	(100.0)				
記載なし															
(%)															
総計	820	4,165	9,717	11,727	9,813	6,045	3,124	1,277	524	278	47,490	20,884	68,374	108.05	33.15
(%)	(1.7)	(8.8)	(20.5)	(24.7)	(20.7)	(12.7)	(6.6)	(2.7)	(1.1)	(0.6)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。

##D4633##

(C)Japanese Society for Dialysis Therapy

2016年調査