

表 1306 Non-HDL-C濃度(mg/dL) 年齢(歳)別 (腹膜透析患者)

年齢	50未満	50 ~	70 ~	90 ~	110 ~	130 ~	150 ~	170 ~	190 ~	210 ~	合計	記載なし	総計	平均	標準偏差
15歳未満				1	3	2	4	1		1	12	73	85	145.42	34.06
(%)				(8.3)	(25.0)	(16.7)	(33.3)	(8.3)		(8.3)	(100.0)				
15歳 ~	1		5	5	5	2	4	1		1	24	70	94	121.96	51.59
(%)	(4.2)		(20.8)	(20.8)	(20.8)	(8.3)	(16.7)	(4.2)		(4.2)	(100.0)				
30歳 ~	1	11	30	47	63	44	21	11	8	4	240	368	608	121.97	34.87
(%)	(0.4)	(4.6)	(12.5)	(19.6)	(26.3)	(18.3)	(8.8)	(4.6)	(3.3)	(1.7)	(100.0)				
45歳 ~	6	33	101	189	220	180	114	75	29	21	968	1,315	2,283	127.06	38.40
(%)	(0.6)	(3.4)	(10.4)	(19.5)	(22.7)	(18.6)	(11.8)	(7.7)	(3.0)	(2.2)	(100.0)				
60歳 ~	9	56	179	314	332	257	189	114	41	43	1,534	2,184	3,718	126.09	39.02
(%)	(0.6)	(3.7)	(11.7)	(20.5)	(21.6)	(16.8)	(12.3)	(7.4)	(2.7)	(2.8)	(100.0)				
75歳 ~	7	22	68	145	145	123	71	50	11	22	664	1,037	1,701	125.90	38.86
(%)	(1.1)	(3.3)	(10.2)	(21.8)	(21.8)	(18.5)	(10.7)	(7.5)	(1.7)	(3.3)	(100.0)				
90歳 ~			7	6	8	17	4	3			45	81	126	126.33	30.08
(%)			(15.6)	(13.3)	(17.8)	(37.8)	(8.9)	(6.7)			(100.0)				
合計	24	122	390	707	776	625	407	255	89	92	3,487	5,128	8,615	126.08	38.53
(%)	(0.7)	(3.5)	(11.2)	(20.3)	(22.3)	(17.9)	(11.7)	(7.3)	(2.6)	(2.6)	(100.0)				
不明															
(%)															
記載なし															
(%)															
総計	24	122	390	707	776	625	407	255	89	92	3,487	5,128	8,615	126.08	38.53
(%)	(0.7)	(3.5)	(11.2)	(20.3)	(22.3)	(17.9)	(11.7)	(7.3)	(2.6)	(2.6)	(100.0)				
平均	63.63	63.51	63.15	63.83	62.79	63.93	63.31	63.99	60.81	64.25	63.41	62.98	63.16		
標準偏差	14.50	12.36	13.46	12.87	13.32	13.61	13.52	12.84	12.49	13.82	13.25	15.26	14.48		

数値下のかっこ内は行方向の合計に対する%です。

##D4641##

(C)Japanese Society for Dialysis Therapy

2016年調査