

表 1307 Non-HDL-C濃度(mg/dL) 透析歴(年)別 (腹膜透析患者)

透析歴	50未満	50~	70~	90~	110~	130~	150~	170~	190~	210~	合計	記載なし	総計	平均	標準偏差
2年未満	7	51	157	295	293	238	169	106	37	35	1,388	2,230	3,618	125.86	37.65
(%)	(0.5)	(3.7)	(11.3)	(21.3)	(21.1)	(17.1)	(12.2)	(7.6)	(2.7)	(2.5)	(100.0)				
2年~	12	34	134	238	270	218	137	86	36	33	1,198	1,786	2,984	126.66	39.27
(%)	(1.0)	(2.8)	(11.2)	(19.9)	(22.5)	(18.2)	(11.4)	(7.2)	(3.0)	(2.8)	(100.0)				
5年~	4	27	80	140	162	129	80	44	13	12	691	884	1,575	124.10	37.13
(%)	(0.6)	(3.9)	(11.6)	(20.3)	(23.4)	(18.7)	(11.6)	(6.4)	(1.9)	(1.7)	(100.0)				
10年~		7	11	26	41	24	14	11	2	6	142	147	289	128.60	41.97
(%)		(4.9)	(7.7)	(18.3)	(28.9)	(16.9)	(9.9)	(7.7)	(1.4)	(4.2)	(100.0)				
15年~		2	6	4	6	11	5	4	1	4	43	42	85	136.72	48.77
(%)		(4.7)	(14.0)	(9.3)	(14.0)	(25.6)	(11.6)	(9.3)	(2.3)	(9.3)	(100.0)				
20年~	1	1	1	4	2	2	1	1		1	14	14	28	117.71	45.28
(%)	(7.1)	(7.1)	(7.1)	(28.6)	(14.3)	(14.3)	(7.1)	(7.1)		(7.1)	(100.0)				
25年~			1		1	1					3	6	9	111.00	29.61
(%)			(33.3)		(33.3)	(33.3)					(100.0)				
30年~					1	1	1	2			4	2	6	161.50	16.42
(%)					(100.0)	(25.0)	(25.0)	(50.0)			(100.0)				
35年~					1						1		1	119.00	
(%)					(100.0)						(100.0)				
40年~								1			1	1	2	176.00	
(%)								(100.0)			(100.0)				
合計	24	122	390	707	776	624	407	255	89	91	3,485	5,112	8,597	126.04	38.48
(%)	(0.7)	(3.5)	(11.2)	(20.3)	(22.3)	(17.9)	(11.7)	(7.3)	(2.6)	(2.6)	(100.0)				
不明						1				1	2	16	18	197.00	79.20
(%)						(50.0)				(50.0)	(100.0)				
記載なし															
(%)															
総計	24	122	390	707	776	625	407	255	89	92	3,487	5,128	8,615	126.08	38.53
(%)	(0.7)	(3.5)	(11.2)	(20.3)	(22.3)	(17.9)	(11.7)	(7.3)	(2.6)	(2.6)	(100.0)				
平均	3.46	3.63	3.19	3.08	3.34	3.41	3.12	3.52	2.81	3.79	3.28	2.85	3.03		
標準偏差	4.25	4.09	3.59	3.37	3.70	3.88	3.60	4.94	3.19	4.49	3.78	3.32	3.52		

数値下のかっこ内は行方向の合計に対する%です。

##D4642##

(C)Japanese Society for Dialysis Therapy

2016年調査