

表 1311 透析前収縮期血圧(mmHg) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差
施設血液透析	32	317	2,857	15,163	45,329	68,505	49,199	18,696	4,574	926	205,598	28,106	233,704	151.05	24.17
(%)	(0.0)	(0.2)	(1.4)	(7.4)	(22.0)	(33.3)	(23.9)	(9.1)	(2.2)	(0.5)	(100.0)				
Off-line HDF	2	18	127	442	951	1,297	849	333	78	21	4,118	519	4,637	146.89	26.11
(%)	(0.0)	(0.4)	(3.1)	(10.7)	(23.1)	(31.5)	(20.6)	(8.1)	(1.9)	(0.5)	(100.0)				
On-line HDF	6	114	781	3,911	11,373	17,223	12,398	4,873	1,254	290	52,223	6,893	59,116	151.32	24.69
(%)	(0.0)	(0.2)	(1.5)	(7.5)	(21.8)	(33.0)	(23.7)	(9.3)	(2.4)	(0.6)	(100.0)				
Push/Pull HDF		1	3	12	18	21	7	2		64	2	66	155.11	22.77	
(%)		(1.6)	(4.7)	(18.8)	(28.1)	(32.8)	(10.9)	(3.1)		(100.0)					
ハイオフィルトレーション		8	16	39	53	55	42	12	6	1	232	20	252	138.53	31.55
(%)		(3.4)	(6.9)	(16.8)	(22.8)	(23.7)	(18.1)	(5.2)	(2.6)	(0.4)	(100.0)				
IHDF	2	27	161	737	2,062	3,137	2,502	1,071	258	51	10,008	720	10,728	152.33	25.15
(%)	(0.0)	(0.3)	(1.6)	(7.4)	(20.6)	(31.3)	(25.0)	(10.7)	(2.6)	(0.5)	(100.0)				
血液濾過			3	3	5	2	5	1		19			19	153.37	31.39
(%)			(15.8)	(15.8)	(26.3)	(10.5)	(26.3)	(5.3)		(100.0)					
血液吸着透析 (リクセル等使用)		8	43	164	308	425	208	67	15	3	1,241	114	1,355	142.63	25.10
(%)		(0.6)	(3.5)	(13.2)	(24.8)	(34.2)	(16.8)	(5.4)	(1.2)	(0.2)	(100.0)				
在宅血液透析			19	69	149	118	61	18	2		436	181	617	138.24	23.31
(%)			(4.4)	(15.8)	(34.2)	(27.1)	(14.0)	(4.1)	(0.5)		(100.0)				
腹膜透析 (CAPD)	2	11	80	399	1,019	986	448	121	19	5	3,090	1,883	4,973	140.64	22.58
(%)	(0.1)	(0.4)	(2.6)	(12.9)	(33.0)	(31.9)	(14.5)	(3.9)	(0.6)	(0.2)	(100.0)				
腹膜透析 (APD)	1	5	48	203	519	541	243	58	12	2	1,632	1,005	2,637	140.70	22.75
(%)	(0.1)	(0.3)	(2.9)	(12.4)	(31.8)	(33.1)	(14.9)	(3.6)	(0.7)	(0.1)	(100.0)				
腹膜透析 (CCPD)	1	1	11	56	195	229	118	39	9	2	661	344	1,005	145.73	22.92
(%)	(0.2)	(0.2)	(1.7)	(8.5)	(29.5)	(34.6)	(17.9)	(5.9)	(1.4)	(0.3)	(100.0)				
合計	46	509	4,144	21,189	61,973	92,539	66,091	25,300	6,230	1,301	279,322	39,787	319,109	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				
記載なし															
(%)															
総計	46	509	4,144	21,189	61,973	92,539	66,091	25,300	6,230	1,301	279,322	39,787	319,109	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4711##

(C)Japanese Society for Dialysis Therapy

2016年調査