

表 1314 透析前収縮期血圧(mmHg) 年齢(歳)別 (透析患者全体)

年齢	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
15歳未満			20	26	11	6	3	1			67	26	93	114.06	22.98
(%)			(29.9)	(38.8)	(16.4)	(9.0)	(4.5)	(1.5)			(100.0)				
15歳～		1	11	81	221	252	142	46	9	5	768	143	911	145.71	23.84
(%)		(0.1)	(1.4)	(10.5)	(28.8)	(32.8)	(18.5)	(6.0)	(1.2)	(0.7)	(100.0)				
30歳～	6	13	155	837	2,491	3,698	2,807	1,226	338	89	11,660	1,762	13,422	152.62	25.26
(%)	(0.1)	(0.1)	(1.3)	(7.2)	(21.4)	(31.7)	(24.1)	(10.5)	(2.9)	(0.8)	(100.0)				
45歳～	9	99	611	3,139	9,952	16,319	12,726	5,423	1,472	363	50,113	7,231	57,344	153.60	24.89
(%)	(0.0)	(0.2)	(1.2)	(6.3)	(19.9)	(32.6)	(25.4)	(10.8)	(2.9)	(0.7)	(100.0)				
60歳～	16	198	1,653	8,482	26,704	41,828	30,368	11,512	2,844	561	124,166	17,463	141,629	151.65	23.99
(%)	(0.0)	(0.2)	(1.3)	(6.8)	(21.5)	(33.7)	(24.5)	(9.3)	(2.3)	(0.5)	(100.0)				
75歳～	15	179	1,533	7,948	21,264	28,764	19,024	6,750	1,491	270	87,238	12,375	99,613	148.29	24.11
(%)	(0.0)	(0.2)	(1.8)	(9.1)	(24.4)	(33.0)	(21.8)	(7.7)	(1.7)	(0.3)	(100.0)				
90歳～		19	161	676	1,329	1,672	1,021	342	76	13	5,309	786	6,095	144.50	25.22
(%)		(0.4)	(3.0)	(12.7)	(25.0)	(31.5)	(19.2)	(6.4)	(1.4)	(0.2)	(100.0)				
合計	46	509	4,144	21,189	61,972	92,539	66,091	25,300	6,230	1,301	279,321	39,786	319,107	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				
不明															
(%)															
記載なし					1						1	1	2	128.00	
(%)					(100.0)						(100.0)				
総計	46	509	4,144	21,189	61,973	92,539	66,091	25,300	6,230	1,301	279,322	39,787	319,109	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				
平均	65.54	69.83	70.00	69.94	69.00	68.21	67.49	66.68	65.69	63.86	68.16	68.03	68.15		
標準偏差	13.98	12.56	13.63	13.05	12.57	12.26	12.21	12.31	12.36	12.69	12.45	12.72	12.48		

数値下のカッコ内は行方向の合計に対する%です。

##D4714##

(C)Japanese Society for Dialysis Therapy

2016年調査