

表 1316 透析前収縮期血圧(mmHg) 原疾患別 (透析患者全体)

原疾患	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	17	196	1,439	7,014	20,118	27,957	17,171	5,364	1,044	185	80,505	11,278	91,783	147.36	23.22
(%)	(0.0)	(0.2)	(1.8)	(8.7)	(25.0)	(34.7)	(21.3)	(6.7)	(1.3)	(0.2)	(100.0)				
慢性腎盂腎炎	17	72	282	599	824	538	188	34	7	2,561	335	2,896	145.92	25.16	
(%)	(0.7)	(2.8)	(11.0)	(23.4)	(32.2)	(21.0)	(7.3)	(1.3)	(0.3)	(100.0)					
急速進行性糸球体腎炎	1	3	44	221	572	766	456	138	26	4	2,231	307	2,538	146.12	23.19
(%)	(0.0)	(0.1)	(2.0)	(9.9)	(25.6)	(34.3)	(20.4)	(6.2)	(1.2)	(0.2)	(100.0)				
妊娠腎/妊娠中毒症	2	6	30	134	343	438	266	93	9	1	1,322	161	1,483	145.69	23.80
(%)	(0.2)	(0.5)	(2.3)	(10.1)	(25.9)	(33.1)	(20.1)	(7.0)	(0.7)	(0.1)	(100.0)				
その他の分類不能の腎炎	1	3	27	146	325	426	256	78	18	2	1,282	172	1,454	145.46	23.90
(%)	(0.1)	(0.2)	(2.1)	(11.4)	(25.4)	(33.2)	(20.0)	(6.1)	(1.4)	(0.2)	(100.0)				
多発性囊胞腎	2	23	168	944	2,877	3,649	1,928	486	79	6	10,162	1,309	11,471	144.96	21.74
(%)	(0.0)	(0.2)	(1.7)	(9.3)	(28.3)	(35.9)	(19.0)	(4.8)	(0.8)	(0.1)	(100.0)				
腎硬化症	3	52	436	2,449	6,886	9,518	5,959	2,012	424	66	27,805	3,845	31,650	148.02	23.46
(%)	(0.0)	(0.2)	(1.6)	(8.8)	(24.8)	(34.2)	(21.4)	(7.2)	(1.5)	(0.2)	(100.0)				
悪性高血圧	4	25	140	545	787	579	220	45	8	2,353	368	2,721	151.41	23.39	
(%)	(0.2)	(1.1)	(5.9)	(23.2)	(33.4)	(24.6)	(9.3)	(1.9)	(0.3)	(100.0)					
糖尿病性腎症	8	93	927	5,666	19,064	34,209	29,855	13,736	3,926	904	108,388	15,269	123,657	156.53	24.63
(%)	(0.0)	(0.1)	(0.9)	(5.2)	(17.6)	(31.6)	(27.5)	(12.7)	(3.6)	(0.8)	(100.0)				
SLE腎炎	3	6	50	181	537	626	374	121	23	7	1,928	294	2,222	145.41	24.39
(%)	(0.2)	(0.3)	(2.6)	(9.4)	(27.9)	(32.5)	(19.4)	(6.3)	(1.2)	(0.4)	(100.0)				
アミロイド腎	3	12	51	106	110	76	21	3	3	382	56	438	142.71	24.51	
(%)	(0.8)	(3.1)	(13.4)	(27.7)	(28.8)	(19.9)	(5.5)	(0.8)		(100.0)					
痛風腎	4	20	104	222	295	199	68	6	6	924	132	1,056	146.66	24.86	
(%)	(0.4)	(2.2)	(11.3)	(24.0)	(31.9)	(21.5)	(7.4)	(0.6)	(0.6)	(100.0)					
先天性代謝異常による腎不全	2	9	38	69	74	49	6	2	2	251	30	281	140.67	25.37	
(%)	(0.8)	(3.6)	(15.1)	(27.5)	(29.5)	(19.5)	(2.4)	(0.8)	(0.8)	(100.0)					
腎・尿路結核	2	7	18	30	41	35	8	4	1	146	22	168	145.64	27.94	
(%)	(1.4)	(4.8)	(12.3)	(20.5)	(28.1)	(24.0)	(5.5)	(2.7)	(0.7)	(100.0)					
腎・尿路結石	2	12	56	133	165	104	29	13		514	64	578	146.10	24.75	
(%)	(0.4)	(2.3)	(10.9)	(25.9)	(32.1)	(20.2)	(5.6)	(2.5)		(100.0)					
腎・尿路腫瘍	14	80	216	289	180	48	20			847	112	959	147.62	23.01	
(%)	(1.7)	(9.4)	(25.5)	(34.1)	(21.3)	(5.7)	(2.4)			(100.0)					
閉塞性尿路障害	2	13	67	170	206	140	41	13	3	655	88	743	146.86	24.69	
(%)	(0.3)	(2.0)	(10.2)	(26.0)	(31.5)	(21.4)	(6.3)	(2.0)	(0.5)	(100.0)					
骨髄腫	1	11	54	76	86	39	11	1		279	44	323	137.89	23.45	
(%)	(0.4)	(3.9)	(19.4)	(27.2)	(30.8)	(14.0)	(3.9)	(0.4)		(100.0)					
腎形成不全	1	20	79	171	180	102	21	8	1	583	81	664	141.92	24.18	
(%)	(0.2)	(3.4)	(13.6)	(29.3)	(30.9)	(17.5)	(3.6)	(1.4)	(0.2)	(100.0)					
不明	3	59	548	2,465	6,610	8,879	5,975	1,979	412	75	27,005	4,407	31,412	147.83	23.99
(%)	(0.0)	(0.2)	(2.0)	(9.1)	(24.5)	(32.9)	(22.1)	(7.3)	(1.5)	(0.3)	(100.0)				
再導入	1	4	55	196	476	634	367	111	26	7	1,877	297	2,174	145.52	24.65
(%)	(0.1)	(0.2)	(2.9)	(10.4)	(25.4)	(33.8)	(19.6)	(5.9)	(1.4)	(0.4)	(100.0)				
その他	4	27	205	804	1,828	2,380	1,443	521	94	16	7,322	1,116	8,438	145.72	24.79
(%)	(0.1)	(0.4)	(2.8)	(11.0)	(25.0)	(32.5)	(19.7)	(7.1)	(1.3)	(0.2)	(100.0)				
合計	46	509	4,144	21,189	61,973	92,539	66,091	25,300	6,230	1,301	279,322	39,787	319,109	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				
記載なし															
総計	46	509	4,144	21,189	61,973	92,539	66,091	25,300	6,230	1,301	279,322	39,787	319,109	150.83	24.37
(%)	(0.0)	(0.2)	(1.5)	(7.6)	(22.2)	(33.1)	(23.7)	(9.1)	(2.2)	(0.5)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4716##

(C)Japanese Society for Dialysis Therapy

2016年調査