

表 1319 透析前収縮期血圧(mmHg) 年齢(歳)別 (施設血液透析患者 透析回数週3回)

年齢	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
15歳未満			1	1		1	1	1			5		5	142.20	34.51
(%)			(20.0)	(20.0)		(20.0)	(20.0)	(20.0)			(100.0)				
15歳～			5	33	131	148	98	28	8	4	455	41	496	148.57	23.48
(%)			(1.1)	(7.3)	(28.8)	(32.5)	(21.5)	(6.2)	(1.8)	(0.9)	(100.0)				
30歳～	4	8	92	507	1,543	2,326	1,794	790	218	51	7,333	481	7,814	152.95	25.15
(%)	(0.1)	(0.1)	(1.3)	(6.9)	(21.0)	(31.7)	(24.5)	(10.8)	(3.0)	(0.7)	(100.0)				
45歳～	8	58	372	1,888	6,204	10,575	8,453	3,624	976	244	32,402	2,090	34,492	154.31	24.73
(%)	(0.0)	(0.2)	(1.1)	(5.8)	(19.1)	(32.6)	(26.1)	(11.2)	(3.0)	(0.8)	(100.0)				
60歳～	11	108	1,019	5,621	18,389	29,522	21,724	8,176	2,057	393	87,020	5,362	92,382	152.16	23.74
(%)	(0.0)	(0.1)	(1.2)	(6.5)	(21.1)	(33.9)	(25.0)	(9.4)	(2.4)	(0.5)	(100.0)				
75歳～	8	115	1,099	5,841	16,063	21,920	14,522	5,128	1,090	201	65,987	4,037	70,024	148.47	23.88
(%)	(0.0)	(0.2)	(1.7)	(8.9)	(24.3)	(33.2)	(22.0)	(7.8)	(1.7)	(0.3)	(100.0)				
90歳～		13	103	507	1,014	1,316	810	261	54	10	4,088	233	4,321	145.12	24.82
(%)		(0.3)	(2.5)	(12.4)	(24.8)	(32.2)	(19.8)	(6.4)	(1.3)	(0.2)	(100.0)				
合計	31	302	2,691	14,398	43,344	65,808	47,402	18,008	4,403	903	197,290	12,244	209,534	151.15	24.14
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.4)	(24.0)	(9.1)	(2.2)	(0.5)	(100.0)				
不明															
(%)															
記載なし					1						1	1	2	128.00	
(%)					(100.0)						(100.0)				
総計	31	302	2,691	14,398	43,345	65,808	47,402	18,008	4,403	903	197,291	12,245	209,536	151.15	24.14
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.4)	(24.0)	(9.1)	(2.2)	(0.5)	(100.0)				
平均	63.68	70.63	71.16	71.03	69.90	69.00	68.18	67.31	66.18	64.40	68.94	68.68	68.93		
標準偏差	13.30	12.54	12.74	12.60	12.35	12.07	12.07	12.19	12.27	12.55	12.25	12.39	12.26		

数値下のカッコ内は行方向の合計に対する%です。

##D4721##

(C)Japanese Society for Dialysis Therapy

2016年調査