

表 1322 透析前収縮期血圧(mmHg) 医療機関別 (施設血液透析患者 透析回数週3回)

医療機関	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差
国公立大学		1	20	51	112	135	69	20	7	2	417	15	432	142.75	25.41
(%)		(0.2)	(4.8)	(12.2)	(26.9)	(32.4)	(16.5)	(4.8)	(1.7)	(0.5)	(100.0)				
私立大学		5	27	101	308	440	319	140	21	4	1,365	192	1,557	150.47	24.67
(%)		(0.4)	(2.0)	(7.4)	(22.6)	(32.2)	(23.4)	(10.3)	(1.5)	(0.3)	(100.0)				
国立病院機構等		1	12	43	82	118	72	31	14	3	376	80	456	147.87	27.19
(%)		(0.3)	(3.2)	(11.4)	(21.8)	(31.4)	(19.1)	(8.2)	(3.7)	(0.8)	(100.0)				
県市町村立・国保	1	19	181	889	2,480	3,951	2,648	982	267	39	11,457	1,066	12,523	150.52	24.12
(%)	(0.0)	(0.2)	(1.6)	(7.8)	(21.6)	(34.5)	(23.1)	(8.6)	(2.3)	(0.3)	(100.0)				
JCHO(旧社会保険)等	1	3	27	169	493	861	578	204	46	9	2,391	56	2,447	151.41	23.41
(%)	(0.0)	(0.1)	(1.1)	(7.1)	(20.6)	(36.0)	(24.2)	(8.5)	(1.9)	(0.4)	(100.0)				
厚生連等		7	74	449	1,327	1,979	1,423	602	150	31	6,042	360	6,402	151.72	24.44
(%)		(0.1)	(1.2)	(7.4)	(22.0)	(32.8)	(23.6)	(10.0)	(2.5)	(0.5)	(100.0)				
その他公的		12	113	515	1,438	1,975	1,458	525	123	22	6,181	362	6,543	149.71	24.46
(%)		(0.2)	(1.8)	(8.3)	(23.3)	(32.0)	(23.6)	(8.5)	(2.0)	(0.4)	(100.0)				
私立総合	1	7	54	374	1,055	1,494	1,151	435	90	23	4,684	174	4,858	151.07	24.07
(%)	(0.0)	(0.1)	(1.2)	(8.0)	(22.5)	(31.9)	(24.6)	(9.3)	(1.9)	(0.5)	(100.0)				
私立	11	110	964	4,947	13,432	19,349	13,705	5,198	1,266	244	59,226	2,858	62,084	150.02	24.54
(%)	(0.0)	(0.2)	(1.6)	(8.4)	(22.7)	(32.7)	(23.1)	(8.8)	(2.1)	(0.4)	(100.0)				
私立診療所	17	137	1,219	6,860	22,618	35,506	25,979	9,871	2,419	526	105,152	7,082	112,234	151.96	23.83
(%)	(0.0)	(0.1)	(1.2)	(6.5)	(21.5)	(33.8)	(24.7)	(9.4)	(2.3)	(0.5)	(100.0)				
総計	31	302	2,691	14,398	43,345	65,808	47,402	18,008	4,403	903	197,291	12,245	209,536	151.15	24.14
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(22.0)	(33.4)	(24.0)	(9.1)	(2.2)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4724##

(C)Japanese Society for Dialysis Therapy

2016年調査