

表 1326 透析前収縮期血圧(mmHg) 原疾患別 (血液透析濾過患者 透析回数週3回)

原疾患	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎 (%)	3 (0.0)	74 (0.3)	428 (2.0)	1,936 (9.1)	5,244 (24.6)	7,350 (34.5)	4,451 (20.9)	1,486 (7.0)	266 (1.2)	50 (0.2)	21,288 (100.0)	1,100	22,388	147.14	23.65
慢性腎盂腎炎 (%)	5 (0.9)	17 (2.9)	63 (10.8)	134 (22.9)	193 (32.9)	120 (20.5)	42 (7.2)	10 (1.7)	2 (0.3)	586 (100.0)	41	627	146.38	25.90	
急速進行性糸球体腎炎 (%)	2 (0.4)	12 (2.5)	46 (9.8)	117 (24.8)	161 (34.2)	97 (20.6)	30 (6.4)	6 (1.3)	471 (100.0)	13	484	145.27	23.59		
妊娠腎/妊娠中毒症 (%)	1 (0.2)	3 (0.7)	12 (2.8)	49 (11.5)	102 (23.9)	145 (34.0)	81 (19.0)	30 (7.0)	4 (0.9)	427 (100.0)	32	459	145.00	24.52	
その他の分類不能の腎炎 (%)			9 (2.7)	39 (11.7)	89 (26.8)	110 (33.1)	60 (18.1)	18 (5.4)	7 (2.1)	332 (100.0)	23	355	145.15	24.37	
多発性囊胞腎 (%)	1 (0.0)	5 (0.2)	43 (1.7)	239 (9.7)	709 (28.8)	849 (34.5)	483 (19.6)	114 (4.6)	18 (0.7)	1 (0.0)	2,462 (100.0)	107	2,569	144.73	21.76
腎硬化症 (%)	15 (0.3)	90 (1.6)	86 (8.6)	482 (23.3)	1,312 (33.4)	1,880 (23.1)	1,299 (7.4)	417 (1.9)	109 (0.3)	17 (0.3)	5,621 (100.0)	275	5,896	149.12	23.91
悪性高血圧 (%)	3 (0.6)	7 (1.3)	29 (5.5)	130 (24.4)	181 (34.0)	127 (23.9)	45 (8.5)	9 (1.7)	1 (0.2)	1 (0.2)	532 (100.0)	23	555	150.06	23.09
糖尿病性腎症 (%)	2 (0.0)	21 (0.1)	200 (0.9)	1,157 (5.0)	3,812 (16.3)	7,110 (30.5)	6,560 (28.1)	3,251 (13.9)	978 (4.2)	253 (1.1)	23,344 (100.0)	1,172	24,516	158.17	25.19
SLE腎炎 (%)	2 (0.4)	4 (0.8)	15 (3.0)	55 (10.9)	133 (26.3)	160 (31.6)	101 (20.0)	26 (5.1)	8 (1.6)	2 (0.4)	506 (100.0)	31	537	144.01	25.86
アミロイド腎 (%)			6 (6.5)	14 (15.2)	26 (28.3)	27 (29.3)	11 (12.0)	7 (7.6)	1 (1.1)	1 (1.1)	92 (100.0)	6	98	140.22	25.98
痛風腎 (%)	1 (0.5)	2 (0.9)	24 (11.3)	55 (25.9)	58 (27.4)	58 (27.4)	12 (27.4)	1 (5.7)	1 (0.5)	1 (0.5)	212 (100.0)	15	227	146.79	23.99
先天性代謝異常による腎不全 (%)			4 (5.7)	7 (10.0)	17 (24.3)	25 (35.7)	12 (17.1)	3 (4.3)	1 (1.4)	1 (1.4)	70 (100.0)	3	73	143.14	26.00
腎・尿路結核 (%)	1 (2.4)	5 (12.2)	5 (12.2)	5 (12.2)	11 (26.8)	9 (22.0)	3 (7.3)	2 (4.9)		41 (100.0)	3	44	143.73	32.97	
腎・尿路結石 (%)			3 (2.8)	10 (9.2)	28 (25.7)	31 (28.4)	27 (24.8)	9 (8.3)	1 (0.9)	1 (0.9)	109 (100.0)	4	113	147.43	24.46
腎・尿路腫瘍 (%)			5 (2.8)	25 (13.9)	46 (25.6)	59 (32.8)	34 (18.9)	7 (3.9)	4 (2.2)	4 (2.2)	180 (100.0)	10	190	144.21	23.54
閉塞性尿路障害 (%)			3 (2.1)	12 (8.2)	40 (27.4)	46 (31.5)	27 (18.5)	14 (9.6)	3 (2.1)	1 (0.7)	146 (100.0)	5	151	148.90	25.75
骨髄腫 (%)			4 (11.1)	11 (30.6)	2 (5.6)	16 (44.4)	2 (5.6)	1 (2.8)		36 (100.0)	1	37	130.25	26.05	
腎形成不全 (%)	1 (0.6)	3 (1.8)	20 (12.1)	48 (29.1)	53 (32.1)	32 (19.4)	7 (4.2)	1 (0.6)		165 (100.0)	6	171	143.75	23.91	
不明 (%)	20 (0.3)	118 (1.9)	525 (8.6)	1,447 (23.8)	1,966 (32.3)	1,391 (22.9)	492 (8.1)	103 (1.7)	18 (0.3)	6,080 (100.0)	451	6,531	148.80	24.24	
再導入 (%)	2 (0.3)	19 (3.1)	64 (10.6)	150 (24.8)	215 (35.5)	111 (18.3)	36 (5.9)	6 (1.0)	3 (0.5)	606 (100.0)	27	633	144.84	24.75	
その他 (%)	1 (0.1)	7 (0.4)	38 (2.3)	180 (10.9)	395 (23.9)	556 (33.7)	336 (20.4)	109 (6.6)	5 (1.5)	1,651 (100.0)	69	1,720	146.51	24.79	
合計 (%)	10 (0.0)	164 (0.3)	1,043 (1.6)	4,992 (7.7)	14,041 (21.6)	21,202 (32.6)	15,429 (23.8)	6,159 (9.5)	1,562 (2.4)	355 (0.5)	64,957 (100.0)	3,417	68,374	151.22	24.90
記載なし (%)															
総計 (%)	10 (0.0)	164 (0.3)	1,043 (1.6)	4,992 (7.7)	14,041 (21.6)	21,202 (32.6)	15,429 (23.8)	6,159 (9.5)	1,562 (2.4)	355 (0.5)	64,957 (100.0)	3,417	68,374	151.22	24.90

数値下のかっこ内は行方向の合計に対する%です。

##D4733#

(C)Japanese Society for Dialysis Therapy

2016年調査