

表 1330 収縮期血圧(mmHg) 透析歴(年)別 (腹膜透析患者)

透析歴	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差
2年未満		7	57	275	726	725	317	71	10	1	2,189	1,429	3,618	140.43	21.77
(%)		(0.3)	(2.6)	(12.6)	(33.2)	(33.1)	(14.5)	(3.2)	(0.5)	(0.0)	(100.0)				
2年~	3	5	46	218	604	597	288	76	16	5	1,858	1,126	2,984	141.67	23.09
(%)	(0.2)	(0.3)	(2.5)	(11.7)	(32.5)	(32.1)	(15.5)	(4.1)	(0.9)	(0.3)	(100.0)				
5年~		2	27	122	316	334	162	53	13	2	1,031	544	1,575	142.68	23.34
(%)		(0.2)	(2.6)	(11.8)	(30.6)	(32.4)	(15.7)	(5.1)	(1.3)	(0.2)	(100.0)				
10年~		2	6	31	55	68	31	12	1	1	207	82	289	141.48	24.80
(%)		(1.0)	(2.9)	(15.0)	(26.6)	(32.9)	(15.0)	(5.8)	(0.5)	(0.5)	(100.0)				
15年~		1	1	4	21	21	7	3			58	27	85	139.95	22.27
(%)		(1.7)	(1.7)	(6.9)	(36.2)	(36.2)	(12.1)	(5.2)			(100.0)				
20年~	1			5	5	6	2	1			20	8	28	133.15	28.13
(%)	(5.0)			(25.0)	(25.0)	(30.0)	(10.0)	(5.0)			(100.0)				
25年~			1		1	1		1			4	5	9	139.75	41.20
(%)			(25.0)		(25.0)	(25.0)		(25.0)			(100.0)				
30年~			1	2			1	1			5	1	6	133.60	45.30
(%)			(20.0)	(40.0)			(20.0)	(20.0)			(100.0)				
35年~											1	1			
(%)															
40年~				1	1						2		2	125.00	18.38
(%)				(50.0)	(50.0)						(100.0)				
合計	4	17	139	658	1,729	1,752	808	218	40	9	5,374	3,223	8,597	141.28	22.74
(%)	(0.1)	(0.3)	(2.6)	(12.2)	(32.2)	(32.6)	(15.0)	(4.1)	(0.7)	(0.2)	(100.0)				
不明					4	4	1				9	9	18	141.00	14.91
(%)					(44.4)	(44.4)	(11.1)				(100.0)				
記載なし															
(%)															
総計	4	17	139	658	1,733	1,756	809	218	40	9	5,383	3,232	8,615	141.28	22.73
(%)	(0.1)	(0.3)	(2.6)	(12.2)	(32.2)	(32.6)	(15.0)	(4.0)	(0.7)	(0.2)	(100.0)				
平均	7.25	4.18	3.34	3.32	3.00	3.10	3.19	3.89	3.68	4.22	3.16	2.80	3.03		
標準偏差	9.18	5.19	4.32	4.20	3.46	3.49	3.43	4.32	2.64	3.07	3.64	3.30	3.52		

数値下のかっこ内は行方向の合計に対する%です。

##D4742##

(C)Japanese Society for Dialysis Therapy

2016年調査