

表 1331 収縮期血圧(mmHg) 原疾患別 (腹膜透析患者)

原疾患	60未満	60~	80~	100~	120~	140~	160~	180~	200~	220~	合計	記載なし	総計	平均	標準偏差	
慢性糸球体腎炎 (%)	2 (0.1)	3 (0.2)	34 (2.0)	200 (11.6)	607 (35.2)	585 (34.0)	219 (12.7)	65 (3.8)	7 (0.4)	1 (0.1)	1,723 (100.0)	943	2,666	140.40	20.95	
慢性腎盂腎炎 (%)		2 (2.9)	1 (1.5)	12 (17.6)	13 (19.1)	24 (35.3)	13 (19.1)	3 (4.4)			68 (100.0)	29	97	140.01	25.55	
急速進行性糸球体腎炎 (%)			2 (4.7)	8 (18.6)	17 (39.5)	12 (27.9)	3 (7.0)	1 (2.3)			43 (100.0)	18	61	134.70	19.77	
妊娠腎/妊娠中毒症 (%)	1 (5.0)			2 (10.0)	4 (20.0)	10 (50.0)	3 (15.0)				20 (100.0)	4	24	140.05	27.17	
その他の分類不能の腎炎 (%)			3 (5.4)	10 (17.9)	17 (30.4)	20 (35.7)	5 (8.9)	1 (1.8)			56 (100.0)	19	75	134.96	21.11	
多発性囊胞腎 (%)	1 (0.6)	5 (3.0)	18 (10.9)	70 (42.4)	50 (30.3)	16 (9.7)	4 (2.4)	1 (0.6)			165 (100.0)	107	272	136.76	20.91	
腎硬化症 (%)	3 (0.4)	16 (2.3)	80 (11.7)	250 (36.7)	223 (32.7)	86 (12.6)	19 (2.8)	5 (0.7)			682 (100.0)	424	1,106	139.38	21.84	
悪性高血圧 (%)		1 (1.6)	6 (9.5)	26 (41.3)	16 (25.4)	10 (15.9)	3 (4.8)	1 (1.6)			63 (100.0)	43	106	142.70	23.27	
糖尿病性腎症 (%)	3 (0.2)	36 (2.1)	178 (10.4)	489 (28.5)	560 (32.7)	328 (19.1)	97 (5.7)	17 (1.0)	6 (0.4)		1,714 (100.0)	1,032	2,746	145.12	23.28	
SLE腎炎 (%)		3 (10.0)	3 (10.0)	7 (23.3)	9 (30.0)	6 (20.0)	1 (3.3)			1 (3.3)	(100.0)	30	55	143.57	29.47	
アミロイド腎 (%)	1 (10.0)	1 (10.0)	2 (20.0)	3 (30.0)	2 (20.0)	1 (10.0)					10 (100.0)	6	16	128.70	27.01	
痛風腎 (%)			2 (16.7)	5 (41.7)	4 (33.3)	1 (8.3)					12 (100.0)	11	23	139.67	18.52	
先天性代謝異常による腎不全 (%)			5 (41.7)	1 (8.3)	3 (25.0)	2 (16.7)	1 (8.3)				12 (100.0)	5	17	137.33	28.07	
腎・尿路結核 (%)												1	1			
腎・尿路結石 (%)				4 (25.0)	7 (43.8)	2 (12.5)	3 (18.8)				16 (100.0)	5	21	133.19	21.49	
腎・尿路腫瘍 (%)					3 (33.3)	4 (44.4)	1 (11.1)			1 (11.1)	(100.0)	9	3	12	150.11	23.97
閉塞性尿路障害 (%)		2 (15.4)	1 (7.7)	5 (38.5)	3 (23.1)	1 (7.7)			1 (7.7)	(100.0)	13	7	20	133.46	31.84	
骨髄腫 (%)	1 (25.0)		2 (50.0)			1 (25.0)				4 (100.0)	3	7	107.75	43.59		
腎形成不全 (%)		9 (20.5)	15 (34.1)	8 (18.2)	7 (15.9)	5 (11.4)				44 (100.0)	21	65	122.75	24.69		
不明 (%)	1 (0.2)	11 (2.2)	69 (13.7)	153 (30.4)	174 (34.5)	73 (14.5)	16 (3.2)	6 (1.2)	1 (0.2)	(100.0)	504	385	889	141.39	22.54	
再導入 (%)		3 (8.1)	5 (13.5)	4 (10.8)	14 (37.8)	8 (21.6)	3 (8.1)			37 (100.0)	28	65	145.57	27.41		
その他 (%)	3 (1.9)	12 (7.6)	36 (22.8)	44 (27.8)	34 (21.5)	24 (15.2)	4 (2.5)	1 (0.6)		158 (100.0)	113	271	132.82	26.62		
合計 (%)	4 (0.1)	17 (0.3)	139 (2.6)	658 (12.2)	1,733 (32.2)	1,756 (32.6)	809 (15.0)	218 (4.0)	40 (0.7)	9 (0.2)	5,383 (100.0)	3,232	8,615	141.28	22.73	
記載なし (%)																
総計 (%)	4 (0.1)	17 (0.3)	139 (2.6)	658 (12.2)	1,733 (32.2)	1,756 (32.6)	809 (15.0)	218 (4.0)	40 (0.7)	9 (0.2)	5,383 (100.0)	3,232	8,615	141.28	22.73	

数値下のかっこ内は行方向の合計に対する%です。

##D4743##

(C)Japanese Society for Dialysis Therapy

2016年調査