

表 1334 透析前拡張期血圧(mmHg) 治療方法(新分類)別(透析患者全体)

| 治療方法 (新分類) | 20未満 | 20~ | 40~ | 60~ | 80~ | 90~ | 100~ | 120~ | 140~ | 合計 | 記載なし | 総計 | 平均 | 標準偏差 |
|---------------------|-------|--------|--------|---------|--------|--------|--------|-------|---------|---------|--------|---------|-------|-------|
| 施設血液透析 | 67 | 718 | 18,450 | 97,111 | 49,292 | 26,221 | 12,559 | 857 | 56 | 205,331 | 28,373 | 233,704 | 77.31 | 14.40 |
| (%) | (0.0) | (0.3) | (9.0) | (47.3) | (24.0) | (12.8) | (6.1) | (0.4) | (0.0) | (100.0) | | | | |
| Off-line HDF | 5 | 25 | 457 | 1,910 | 936 | 529 | 223 | 21 | 1 | 4,107 | 530 | 4,637 | 76.24 | 15.19 |
| (%) | (0.1) | (0.6) | (11.1) | (46.5) | (22.8) | (12.9) | (5.4) | (0.5) | (0.0) | (100.0) | | | | |
| On-line HDF | 16 | 193 | 3,688 | 22,382 | 13,510 | 7,947 | 4,146 | 277 | 25 | 52,184 | 6,932 | 59,116 | 79.53 | 14.66 |
| (%) | (0.0) | (0.4) | (7.1) | (42.9) | (25.9) | (15.2) | (7.9) | (0.5) | (0.0) | (100.0) | | | | |
| Push/Pull HDF | | 4 | 30 | 15 | 9 | 6 | | | 64 | 2 | 66 | 78.66 | 13.97 | |
| (%) | | (6.3) | (46.9) | (23.4) | (14.1) | (9.4) | | | (100.0) | | | | | |
| バイオフィルレーション | 5 | 39 | 114 | 42 | 20 | 10 | | | 230 | 22 | 252 | 72.12 | 16.05 | |
| (%) | (2.2) | (17.0) | (49.6) | (18.3) | (8.7) | (4.3) | | | (100.0) | | | | | |
| IHDF | 5 | 36 | 932 | 4,659 | 2,454 | 1,260 | 611 | 47 | 5 | 10,009 | 719 | 10,728 | 77.42 | 14.58 |
| (%) | (0.0) | (0.4) | (9.3) | (46.5) | (24.5) | (12.6) | (6.1) | (0.5) | (0.0) | (100.0) | | | | |
| 血液濾過 | | 2 | 7 | 6 | 3 | | 1 | | 19 | | 19 | 78.63 | 15.20 | |
| (%) | | (10.5) | (36.8) | (31.6) | (15.8) | | (5.3) | | (100.0) | | | | | |
| 血液吸着透析 (リクセル等使用) | 6 | 155 | 644 | 260 | 115 | 51 | 1 | | 1,232 | 123 | 1,355 | 74.18 | 13.99 | |
| (%) | (0.5) | (12.6) | (52.3) | (21.1) | (9.3) | (4.1) | (0.1) | | (100.0) | | | | | |
| 在宅血液透析 | 1 | 40 | 178 | 118 | 58 | 38 | 4 | | 437 | 180 | 617 | 79.23 | 15.08 | |
| (%) | (0.2) | (9.2) | (40.7) | (27.0) | (13.3) | (8.7) | (0.9) | | (100.0) | | | | | |
| 腹膜透析 (CAPD) | 10 | 239 | 1,340 | 803 | 428 | 231 | 13 | | 3,064 | 1,909 | 4,973 | 78.72 | 14.31 | |
| (%) | (0.3) | (7.8) | (43.7) | (26.2) | (14.0) | (7.5) | (0.4) | | (100.0) | | | | | |
| 腹膜透析 (APD) | 4 | 140 | 652 | 427 | 255 | 133 | 10 | | 1,621 | 1,016 | 2,637 | 79.44 | 14.65 | |
| (%) | (0.2) | (8.6) | (40.2) | (26.3) | (15.7) | (8.2) | (0.6) | | (100.0) | | | | | |
| 腹膜透析 (CCPD) | | 31 | 262 | 182 | 99 | 73 | 6 | | 653 | 352 | 1,005 | 81.80 | 14.36 | |
| (%) | | (4.7) | (40.1) | (27.9) | (15.2) | (11.2) | (0.9) | | (100.0) | | | | | |
| 合計 | 93 | 998 | 24,177 | 129,289 | 68,045 | 36,944 | 18,081 | 1,237 | 87 | 278,951 | 40,158 | 319,109 | 77.73 | 14.50 |
| (%) | (0.0) | (0.4) | (8.7) | (46.3) | (24.4) | (13.2) | (6.5) | (0.4) | (0.0) | (100.0) | | | | |
| 記載なし | | | | | | | | | | | | | | |
| 総計 | 93 | 998 | 24,177 | 129,289 | 68,045 | 36,944 | 18,081 | 1,237 | 87 | 278,951 | 40,158 | 319,109 | 77.73 | 14.50 |
| (%) | (0.0) | (0.4) | (8.7) | (46.3) | (24.4) | (13.2) | (6.5) | (0.4) | (0.0) | (100.0) | | | | |

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4811##

(C)Japanese Society for Dialysis Therapy

2016年調査