

表 1339 透析前拡張期血圧(mmHg) 原疾患別 (透析患者全体)

原疾患	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	21	319	6,175	36,262	20,656	11,319	5,242	336	21	80,351	11,432	91,783	78.32	14.33
(%)	(0.0)	(0.4)	(7.7)	(45.1)	(25.7)	(14.1)	(6.5)	(0.4)	(0.0)	(100.0)				
慢性腎盂腎炎		14	242	1,145	616	350	174	14	1	2,556	340	2,896	77.78	15.03
(%)		(0.5)	(9.5)	(44.8)	(24.1)	(13.7)	(6.8)	(0.5)	(0.0)	(100.0)				
急速進行性糸球体腎炎		5	194	1,070	522	282	150	8		2,231	307	2,538	77.49	14.25
(%)		(0.2)	(8.7)	(48.0)	(23.4)	(12.6)	(6.7)	(0.4)		(100.0)				
妊娠腎/妊娠中毒症	2	8	110	600	342	167	89	2		1,320	163	1,483	77.47	14.59
(%)	(0.2)	(0.6)	(8.3)	(45.5)	(25.9)	(12.7)	(6.7)	(0.2)		(100.0)				
その他の分類不能の腎炎	1	3	104	550	326	184	105	7	1	1,281	173	1,454	78.97	14.90
(%)	(0.1)	(0.2)	(8.1)	(42.9)	(25.4)	(14.4)	(8.2)	(0.5)	(0.1)	(100.0)				
多発性嚢胞腎	1	20	601	4,410	2,830	1,544	713	25	2	10,146	1,325	11,471	79.54	13.52
(%)	(0.0)	(0.2)	(5.9)	(43.5)	(27.9)	(15.2)	(7.0)	(0.2)	(0.0)	(100.0)				
腎硬化症	5	107	2,985	13,746	6,254	3,082	1,498	93	9	27,779	3,871	31,650	75.95	14.33
(%)	(0.0)	(0.4)	(10.7)	(49.5)	(22.5)	(11.1)	(5.4)	(0.3)	(0.0)	(100.0)				
悪性高血圧		8	145	897	569	434	256	39	2	2,350	371	2,721	81.95	16.01
(%)		(0.3)	(6.2)	(38.2)	(24.2)	(18.5)	(10.9)	(1.7)	(0.1)	(100.0)				
糖尿病性腎症	46	317	9,554	50,891	25,808	14,006	7,107	517	31	108,277	15,380	123,657	77.62	14.51
(%)	(0.0)	(0.3)	(8.8)	(47.0)	(23.8)	(12.9)	(6.6)	(0.5)	(0.0)	(100.0)				
SLE腎炎	1	9	150	836	492	281	145	10		1,924	298	2,222	78.67	14.81
(%)	(0.1)	(0.5)	(7.8)	(43.5)	(25.6)	(14.6)	(7.5)	(0.5)		(100.0)				
アミロイド腎		1	50	160	94	53	23		1	382	56	438	76.90	15.39
(%)		(0.3)	(13.1)	(41.9)	(24.6)	(13.9)	(6.0)		(0.3)	(100.0)				
痛風腎		5	67	441	232	116	57	5		923	133	1,056	77.87	14.13
(%)		(0.5)	(7.3)	(47.8)	(25.1)	(12.6)	(6.2)	(0.5)		(100.0)				
先天性代謝異常による腎不全		4	23	88	71	39	22	3	1	251	30	281	80.00	17.49
(%)		(1.6)	(9.2)	(35.1)	(28.3)	(15.5)	(8.8)	(1.2)	(0.4)	(100.0)				
腎・尿路結核		1	16	72	37	15	5			146	22	168	74.23	14.65
(%)		(0.7)	(11.0)	(49.3)	(25.3)	(10.3)	(3.4)			(100.0)				
腎・尿路結石	1	3	46	247	133	58	24	2		514	64	578	76.19	14.15
(%)	(0.2)	(0.6)	(8.9)	(48.1)	(25.9)	(11.3)	(4.7)	(0.4)		(100.0)				
腎・尿路腫瘍			77	394	228	99	47	1		846	113	959	77.02	13.24
(%)			(9.1)	(46.6)	(27.0)	(11.7)	(5.6)	(0.1)		(100.0)				
閉塞性尿路障害	1	3	38	300	164	95	48	4	1	654	89	743	79.22	14.79
(%)	(0.2)	(0.5)	(5.8)	(45.9)	(25.1)	(14.5)	(7.3)	(0.6)	(0.2)	(100.0)				
骨髄腫		1	33	143	75	22	5			279	44	323	74.08	12.80
(%)		(0.4)	(11.8)	(51.3)	(26.9)	(7.9)	(1.8)			(100.0)				
腎形成不全	2	3	41	219	154	100	55	7		581	83	664	80.79	16.05
(%)	(0.3)	(0.5)	(7.1)	(37.7)	(26.5)	(17.2)	(9.5)	(1.2)		(100.0)				
不明	8	113	2,657	12,722	6,284	3,412	1,645	120	10	26,971	4,441	31,412	77.07	14.67
(%)	(0.0)	(0.4)	(9.9)	(47.2)	(23.3)	(12.7)	(6.1)	(0.4)	(0.0)	(100.0)				
再導入	1	8	123	762	489	310	173	9	1	1,876	298	2,174	80.20	15.08
(%)	(0.1)	(0.4)	(6.6)	(40.6)	(26.1)	(16.5)	(9.2)	(0.5)	(0.1)	(100.0)				
その他	3	46	746	3,334	1,669	976	498	35	6	7,313	1,125	8,438	77.30	15.29
(%)	(0.0)	(0.6)	(10.2)	(45.6)	(22.8)	(13.3)	(6.8)	(0.5)	(0.1)	(100.0)				
合計	93	998	24,177	129,289	68,045	36,944	18,081	1,237	87	278,951	40,158	319,109	77.73	14.50
(%)	(0.0)	(0.4)	(8.7)	(46.3)	(24.4)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	93	998	24,177	129,289	68,045	36,944	18,081	1,237	87	278,951	40,158	319,109	77.73	14.50
(%)	(0.0)	(0.4)	(8.7)	(46.3)	(24.4)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

##D4816##

(C)Japanese Society for Dialysis Therapy

2016年調査