

表 1340 透析前拡張期血圧(mmHg) 医療機関別 (透析患者全体)

医療機関	20未満	20~	40~	60~	80~	90~	100~	120~	140~	合計	記載なし	総計	平均	標準偏差
国公立大学	1	5	96	374	183	99	42	2	1	803	321	1,124	75.87	15.23
(%)	(0.1)	(0.6)	(12.0)	(46.6)	(22.8)	(12.3)	(5.2)	(0.2)	(0.1)	(100.0)				
私立大学	2	15	198	1,104	607	319	192	16	2	2,455	1,231	3,686	78.37	15.11
(%)	(0.1)	(0.6)	(8.1)	(45.0)	(24.7)	(13.0)	(7.8)	(0.7)	(0.1)	(100.0)				
国立病院・機構等		2	65	288	153	101	46	3		658	253	911	77.93	14.73
(%)		(0.3)	(9.9)	(43.8)	(23.3)	(15.3)	(7.0)	(0.5)		(100.0)				
県市町村立・国保	7	69	1,448	7,388	3,986	2,189	1,041	83	5	16,216	3,532	19,748	77.86	14.62
(%)	(0.0)	(0.4)	(8.9)	(45.6)	(24.6)	(13.5)	(6.4)	(0.5)	(0.0)	(100.0)				
JCHO(旧社会保険)等	1	11	251	1,472	831	442	228	12	1	3,249	516	3,765	78.54	14.27
(%)	(0.0)	(0.3)	(7.7)	(45.3)	(25.6)	(13.6)	(7.0)	(0.4)	(0.0)	(100.0)				
厚生連等	3	25	777	3,689	1,996	1,132	592	40	4	8,258	1,127	9,385	78.17	14.81
(%)	(0.0)	(0.3)	(9.4)	(44.7)	(24.2)	(13.7)	(7.2)	(0.5)	(0.0)	(100.0)				
その他公的	2	45	802	3,843	2,045	1,136	591	46	2	8,512	2,346	10,858	77.75	14.87
(%)	(0.0)	(0.5)	(9.4)	(45.1)	(24.0)	(13.3)	(6.9)	(0.5)	(0.0)	(100.0)				
私立総合	2	29	602	2,995	1,646	916	445	27	1	6,663	1,150	7,813	77.94	14.60
(%)	(0.0)	(0.4)	(9.0)	(44.9)	(24.7)	(13.7)	(6.7)	(0.4)	(0.0)	(100.0)				
私立	25	379	8,062	38,199	18,920	9,867	4,754	313	17	80,536	11,053	91,589	76.80	14.59
(%)	(0.0)	(0.5)	(10.0)	(47.4)	(23.5)	(12.3)	(5.9)	(0.4)	(0.0)	(100.0)				
私立診療所	50	418	11,876	69,937	37,678	20,743	10,150	695	54	151,601	18,629	170,230	78.17	14.36
(%)	(0.0)	(0.3)	(7.8)	(46.1)	(24.9)	(13.7)	(6.7)	(0.5)	(0.0)	(100.0)				
総計	93	998	24,177	129,289	68,045	36,944	18,081	1,237	87	278,951	40,158	319,109	77.73	14.50
(%)	(0.0)	(0.4)	(8.7)	(46.3)	(24.4)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4817##

(C)Japanese Society for Dialysis Therapy

2016年調査