

表 1357 透析前脈拍(拍/分) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	40未満	40~	50~	60~	70~	80~	90~	100~	110~	120~	合計	記載なし	総計	平均	標準偏差
施設血液透析	147	2,187	17,511	52,549	62,735	39,869	17,021	5,159	1,202	494	198,874	34,830	233,704	74.43	12.73
(%)	(0.1)	(1.1)	(8.8)	(26.4)	(31.5)	(20.0)	(8.6)	(2.6)	(0.6)	(0.2)	(100.0)				
Off-line HDF	2	37	283	1,048	1,185	867	373	143	32	11	3,981	656	4,637	75.53	13.03
(%)	(0.1)	(0.9)	(7.1)	(26.3)	(29.8)	(21.8)	(9.4)	(3.6)	(0.8)	(0.3)	(100.0)				
On-line HDF	38	403	4,000	12,670	15,806	11,069	4,669	1,466	365	163	50,649	8,467	59,116	75.39	12.81
(%)	(0.1)	(0.8)	(7.9)	(25.0)	(31.2)	(21.9)	(9.2)	(2.9)	(0.7)	(0.3)	(100.0)				
Push/Pull HDF		6	10	22	13	4	3		2	60	6	66	77.27	15.79	
(%)		(10.0)	(16.7)	(36.7)	(21.7)	(6.7)	(5.0)		(3.3)	(100.0)					
バイオフィルトレーション		3	13	61	66	51	16	11	4	1	226	26	252	75.82	13.75
(%)		(1.3)	(5.8)	(27.0)	(29.2)	(22.6)	(7.1)	(4.9)	(1.8)	(0.4)	(100.0)				
IHDF	5	124	855	2,575	3,023	1,956	873	264	52	18	9,745	983	10,728	74.39	12.77
(%)	(0.1)	(1.3)	(8.8)	(26.4)	(31.0)	(20.1)	(9.0)	(2.7)	(0.5)	(0.2)	(100.0)				
血液濾過			4	3	2	1	2			12	7	19	79.25	14.12	
(%)			(33.3)	(25.0)	(16.7)	(8.3)	(16.7)			(100.0)					
血液吸着透析 (リクセル等使用)	2	11	88	332	384	242	95	29	14	4	1,201	154	1,355	74.77	12.76
(%)	(0.2)	(0.9)	(7.3)	(27.6)	(32.0)	(20.1)	(7.9)	(2.4)	(1.2)	(0.3)	(100.0)				
在宅血液透析		1	13	78	133	96	49	23	7	3	403	214	617	79.09	13.25
(%)		(0.2)	(3.2)	(19.4)	(33.0)	(23.8)	(12.2)	(5.7)	(1.7)	(0.7)	(100.0)				
腹膜透析 (CAPD)		7	135	527	855	665	342	112	28	8	2,679	2,294	4,973	78.07	12.78
(%)		(0.3)	(5.0)	(19.7)	(31.9)	(24.8)	(12.8)	(4.2)	(1.0)	(0.3)	(100.0)				
腹膜透析 (APD)	1	8	87	272	381	352	181	55	17	14	1,368	1,269	2,637	78.26	13.89
(%)	(0.1)	(0.6)	(6.4)	(19.9)	(27.9)	(25.7)	(13.2)	(4.0)	(1.2)	(1.0)	(100.0)				
腹膜透析 (CCPD)		3	28	126	189	130	75	22	6		579	426	1,005	77.48	12.53
(%)		(0.5)	(4.8)	(21.8)	(32.6)	(22.5)	(13.0)	(3.8)	(1.0)		(100.0)				
合計	195	2,784	23,019	70,252	84,782	55,312	23,699	7,289	1,727	718	269,777	49,332	319,109	74.70	12.77
(%)	(0.1)	(1.0)	(8.5)	(26.0)	(31.4)	(20.5)	(8.8)	(2.7)	(0.6)	(0.3)	(100.0)				
記載なし															
(%)															
総計	195	2,784	23,019	70,252	84,782	55,312	23,699	7,289	1,727	718	269,777	49,332	319,109	74.70	12.77
(%)	(0.1)	(1.0)	(8.5)	(26.0)	(31.4)	(20.5)	(8.8)	(2.7)	(0.6)	(0.3)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

##D4911##

(C)Japanese Society for Dialysis Therapy

2016年調査