

表 1367 透析前脈拍(拍/分) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	40未満	40~	50~	60~	70~	80~	90~	100~	110~	120~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	34	515	4,924	14,831	16,680	9,554	3,817	1,224	290	131	52,000	5,172	57,172	73.62	12.49
(%)	(0.1)	(1.0)	(9.5)	(28.5)	(32.1)	(18.4)	(7.3)	(2.4)	(0.6)	(0.3)	(100.0)				
慢性腎盂腎炎	1	13	135	455	542	356	151	46	13	5	1,717	151	1,868	74.88	12.60
(%)	(0.1)	(0.8)	(7.9)	(26.5)	(31.6)	(20.7)	(8.8)	(2.7)	(0.8)	(0.3)	(100.0)				
急速進行性糸球体腎炎	1	17	114	413	508	311	135	52	13	4	1,568	119	1,687	75.23	12.81
(%)	(0.1)	(1.1)	(7.3)	(26.3)	(32.4)	(19.8)	(8.6)	(3.3)	(0.8)	(0.3)	(100.0)				
妊娠腎/妊娠中毒症	1	6	49	210	282	145	60	21	5	3	782	67	849	74.73	12.34
(%)	(0.1)	(0.8)	(6.3)	(26.9)	(36.1)	(18.5)	(7.7)	(2.7)	(0.6)	(0.4)	(100.0)				
その他の分類不能の腎炎	3	11	59	217	262	173	75	19	10		829	73	902	74.99	12.68
(%)	(0.4)	(1.3)	(7.1)	(26.2)	(31.6)	(20.9)	(9.0)	(2.3)	(1.2)		(100.0)				
多発性囊胞腎	1	71	663	2,033	2,186	1,258	472	149	36	11	6,880	668	7,548	73.26	12.41
(%)	(0.0)	(1.0)	(9.6)	(29.5)	(31.8)	(18.3)	(6.9)	(2.2)	(0.5)	(0.2)	(100.0)				
腎硬化症	20	309	2,379	5,890	5,902	3,390	1,345	402	99	41	19,777	1,741	21,518	72.42	12.68
(%)	(0.1)	(1.6)	(12.0)	(29.8)	(29.8)	(17.1)	(6.8)	(2.0)	(0.5)	(0.2)	(100.0)				
悪性高血圧	11	166	418	476	352	136	40	12	3		1,614	163	1,777	74.41	12.58
(%)	(0.7)	(10.3)	(25.9)	(29.5)	(21.8)	(8.4)	(2.5)	(0.7)	(0.2)		(100.0)				
糖尿病性腎症	53	817	5,723	18,149	24,431	17,143	7,738	2,230	475	185	76,944	7,524	84,468	75.62	12.74
(%)	(0.1)	(1.1)	(7.4)	(23.6)	(31.8)	(22.3)	(10.1)	(2.9)	(0.6)	(0.2)	(100.0)				
SLE腎炎	11	60	308	419	296	113	43	10	4		1,264	106	1,370	76.31	12.43
(%)	(0.9)	(4.7)	(24.4)	(33.1)	(23.4)	(8.9)	(3.4)	(0.8)	(0.3)		(100.0)				
アミロイド腎	1	12	74	81	62	18	8	4			260	20	280	75.74	12.10
(%)	(0.4)	(4.6)	(28.5)	(31.2)	(23.8)	(6.9)	(3.1)	(1.5)			(100.0)				
痛風腎	1	8	64	184	196	115	52	13	3	2	638	62	700	73.45	12.56
(%)	(0.2)	(1.3)	(10.0)	(28.8)	(30.7)	(18.0)	(8.2)	(2.0)	(0.5)	(0.3)	(100.0)				
先天性代謝異常による腎不全	2	10	33	51	34	17	5	3			155	14	169	76.44	13.22
(%)	(1.3)	(6.5)	(21.3)	(32.9)	(21.9)	(11.0)	(3.2)	(1.9)			(100.0)				
腎・尿路結核	13	23	31	20	5	3					95	11	106	72.69	11.73
(%)		(13.7)	(24.2)	(32.6)	(21.1)	(5.3)	(3.2)				(100.0)				
腎・尿路結石	4	28	104	111	67	28	11	4	1		358	26	384	74.26	12.92
(%)	(1.1)	(7.8)	(29.1)	(31.0)	(18.7)	(7.8)	(3.1)	(1.1)	(0.3)		(100.0)				
腎・尿路腫瘍	1	7	60	166	175	119	51	18	3	5	605	39	644	74.08	13.77
(%)	(0.2)	(1.2)	(9.9)	(27.4)	(28.9)	(19.7)	(8.4)	(3.0)	(0.5)	(0.8)	(100.0)				
閉塞性尿路障害	2	35	118	132	95	44	10	2			438	27	465	75.03	12.09
(%)	(0.5)	(8.0)	(26.9)	(30.1)	(21.7)	(10.0)	(2.3)	(0.5)			(100.0)				
骨髄腫	2	24	43	62	43	16	3			1	194	19	213	74.28	12.31
(%)	(1.0)	(12.4)	(22.2)	(32.0)	(22.2)	(8.2)	(1.5)			(0.5)	(100.0)				
腎形成不全	2	19	78	102	82	38	12	3			336	32	368	76.91	12.36
(%)	(0.6)	(5.7)	(23.2)	(30.4)	(24.4)	(11.3)	(3.6)	(0.9)			(100.0)				
不明	15	222	1,775	5,170	5,786	3,480	1,489	456	123	54	18,570	1,908	20,478	73.87	12.80
(%)	(0.1)	(1.2)	(9.6)	(27.8)	(31.2)	(18.7)	(8.0)	(2.5)	(0.7)	(0.3)	(100.0)				
再導入	4	9	74	249	345	232	120	29	13	2	1,077	87	1,164	75.96	12.94
(%)	(0.4)	(0.8)	(6.9)	(23.1)	(32.0)	(21.5)	(11.1)	(2.7)	(1.2)	(0.2)	(100.0)				
その他	4	67	430	1,256	1,552	996	424	155	31	13	4,928	478	5,406	74.68	13.08
(%)	(0.1)	(1.4)	(8.7)	(25.5)	(31.5)	(20.2)	(8.6)	(3.1)	(0.6)	(0.3)	(100.0)				
合計	139	2,107	16,816	50,422	60,312	38,323	16,344	4,949	1,152	465	191,029	18,507	209,536	74.43	12.72
(%)	(0.1)	(1.1)	(8.8)	(26.4)	(31.6)	(20.1)	(8.6)	(2.6)	(0.6)	(0.2)	(100.0)				
記載なし															
総計	139	2,107	16,816	50,422	60,312	38,323	16,344	4,949	1,152	465	191,029	18,507	209,536	74.43	12.72
(%)	(0.1)	(1.1)	(8.8)	(26.4)	(31.6)	(20.1)	(8.6)	(2.6)	(0.6)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

##D4923##

(C)Japanese Society for Dialysis Therapy

2016年調査