

表 1372 透析前脈拍(拍/分) 原疾患別 (血液透析濾過患者 透析回数週3回)

原疾患	40未満	40~	50~	60~	70~	80~	90~	100~	110~	120~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12	177	1,750	5,601	6,507	4,117	1,660	532	140	47	20,543	1,845	22,388	74.43	12.59
(%)	(0.1)	(0.9)	(8.5)	(27.3)	(31.7)	(20.0)	(8.1)	(2.6)	(0.7)	(0.2)	(100.0)				
慢性腎盂腎炎	1	1	46	148	168	129	56	12	2	1	564	63	627	75.10	12.13
(%)	(0.2)	(0.2)	(8.2)	(26.2)	(29.8)	(22.9)	(9.9)	(2.1)	(0.4)	(0.2)	(100.0)				
急速進行性糸球体腎炎	1	2	36	107	133	114	45	17	1	3	459	25	484	76.10	13.16
(%)	(0.2)	(0.4)	(7.8)	(23.3)	(29.0)	(24.8)	(9.8)	(3.7)	(0.2)	(0.7)	(100.0)				
妊娠腎/妊娠中毒症	3	27	118	138	83	37	13	3	1	423	36	459	75.31	12.77	
(%)	(0.7)	(6.4)	(27.9)	(32.6)	(19.6)	(8.7)	(3.1)	(0.7)	(0.2)	(100.0)					
その他の分類不能の腎炎	3	20	83	114	72	20	12	2		326	29	355	75.14	11.84	
(%)	(0.9)	(6.1)	(25.5)	(35.0)	(22.1)	(6.1)	(3.7)	(0.6)		(100.0)					
多発性囊胞腎	1	19	226	681	811	419	162	60	10	7	2,396	173	2,569	73.56	12.17
(%)	(0.0)	(0.8)	(9.4)	(28.4)	(33.8)	(17.5)	(6.8)	(2.5)	(0.4)	(0.3)	(100.0)				
腎硬化症	6	78	637	1,562	1,649	972	370	134	33	13	5,454	442	5,896	72.89	12.80
(%)	(0.1)	(1.4)	(11.7)	(28.6)	(30.2)	(17.8)	(6.8)	(2.5)	(0.6)	(0.2)	(100.0)				
悪性高血圧	6	36	129	160	106	47	26	4	1	515	40	555	75.83	13.14	
(%)	(1.2)	(7.0)	(25.0)	(31.1)	(20.6)	(9.1)	(5.0)	(0.8)	(0.2)	(100.0)					
糖尿病性腎症	21	182	1,520	5,008	6,912	5,469	2,525	763	166	85	22,651	1,865	24,516	76.69	12.98
(%)	(0.1)	(0.8)	(6.7)	(22.1)	(30.5)	(24.1)	(11.1)	(3.4)	(0.7)	(0.4)	(100.0)				
SLE腎炎	4	35	146	134	111	44	14	6		494	43	537	75.29	12.72	
(%)	(0.8)	(7.1)	(29.6)	(27.1)	(22.5)	(8.9)	(2.8)	(1.2)		(100.0)					
アミロイド腎	3	24	32	24	5	4				92	6	98	75.86	11.26	
(%)	(3.3)	(26.1)	(34.8)	(26.1)	(5.4)	(4.3)				(100.0)					
痛風腎	1	17	50	71	43	15	5	6	1	209	18	227	75.89	13.76	
(%)	(0.5)	(8.1)	(23.9)	(34.0)	(20.6)	(7.2)	(2.4)	(2.9)	(0.5)	(100.0)					
先天性代謝異常による腎不全	4	18	18	20	5	3				68	5	73	76.40	11.26	
(%)	(5.9)	(26.5)	(26.5)	(29.4)	(7.4)	(4.4)				(100.0)					
腎・尿路結核	1	2	13	12	7	5				40	4	44	73.95	11.65	
(%)	(2.5)	(5.0)	(32.5)	(30.0)	(17.5)	(12.5)				(100.0)					
腎・尿路結石	1	8	29	36	19	7	2	1		103	10	113	73.99	12.19	
(%)	(1.0)	(7.8)	(28.2)	(35.0)	(18.4)	(6.8)	(1.9)	(1.0)		(100.0)					
腎・尿路腫瘍	1	19	42	59	46	7	2	1		177	13	190	73.09	11.03	
(%)	(0.6)	(10.7)	(23.7)	(33.3)	(26.0)	(4.0)	(1.1)	(0.6)		(100.0)					
閉塞性尿路障害	1	10	39	48	29	12	2	1		142	9	151	74.55	11.62	
(%)	(0.7)	(7.0)	(27.5)	(33.8)	(20.4)	(8.5)	(1.4)	(0.7)		(100.0)					
骨髄腫	2	14	12	4	3					35	2	37	71.46	10.05	
(%)	(5.7)	(40.0)	(34.3)	(11.4)	(8.6)					(100.0)					
腎形成不全	10	29	54	44	17	3	3	1		161	10	171	77.32	12.66	
(%)	(6.2)	(18.0)	(33.5)	(27.3)	(10.6)	(1.9)	(1.9)	(0.6)		(100.0)					
不明	1	50	482	1,550	1,818	1,310	508	171	44	20	5,954	577	6,531	75.14	12.85
(%)	(0.0)	(0.8)	(8.1)	(26.0)	(30.5)	(22.0)	(8.5)	(2.9)	(0.7)	(0.3)	(100.0)				
再導入	2	37	133	197	132	54	21	6	3	585	48	633	76.55	12.83	
(%)	(0.3)	(6.3)	(22.7)	(33.7)	(22.6)	(9.2)	(3.6)	(1.0)	(0.5)	(100.0)					
その他	1	19	98	410	518	326	178	42	11	9	1,612	108	1,720	75.80	13.18
(%)	(0.1)	(1.2)	(6.1)	(25.4)	(32.1)	(20.2)	(11.0)	(2.6)	(0.7)	(0.6)	(100.0)				
合計	44	551	5,025	15,934	19,601	13,596	5,782	1,838	440	192	63,003	5,371	68,374	75.25	12.83
(%)	(0.1)	(0.9)	(8.0)	(25.3)	(31.1)	(21.6)	(9.2)	(2.9)	(0.7)	(0.3)	(100.0)				
記載なし															
総計	44	551	5,025	15,934	19,601	13,596	5,782	1,838	440	192	63,003	5,371	68,374	75.25	12.83
(%)	(0.1)	(0.9)	(8.0)	(25.3)	(31.1)	(21.6)	(9.2)	(2.9)	(0.7)	(0.3)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D4933##

(C)Japanese Society for Dialysis Therapy

2016年調査