

表 1409 使用透析液の種類 原疾患別（腹膜透析患者）

原疾患	1.5%液 のみ (イコデキストリン なし)	1.5%液と 2.5%液の 併用 (イコデキストリン なし)	2.5%液 のみ (イコデキストリン なし)	4.25%液 のみ (イコデキストリン なし)	イコデキストリン 単独 (ブドウ糖 使用なし)	1.5%液+ イコデキストリン	1.5%液+ 2.5%液+ イコデキストリン	2.5%液+ イコデキストリン	4.25%液+ イコデキストリン	合計	不明	記載なし	総計
慢性糸球体腎炎	702	248	46		12	462	223	145		1,838	6	822	2,666
(%)	(38.2)	(13.5)	(2.5)		(0.7)	(25.1)	(12.1)	(7.9)		(100.0)			
慢性腎盂腎炎	30	7	4	2		19	10	3		75		22	97
(%)	(40.0)	(9.3)	(5.3)	(2.7)		(25.3)	(13.3)	(4.0)		(100.0)			
急速進行性糸球体腎炎	11	8	1		1	9	9	4	1	44		17	61
(%)	(25.0)	(18.2)	(2.3)		(2.3)	(20.5)	(20.5)	(9.1)	(2.3)	(100.0)			
妊娠腎/妊娠中毒症	7	3	1				4	5		20		4	24
(%)	(35.0)	(15.0)	(5.0)				(20.0)	(25.0)		(100.0)			
その他の分類不能の腎炎	17	9	1			13	8	2		50		25	75
(%)	(34.0)	(18.0)	(2.0)			(26.0)	(16.0)	(4.0)		(100.0)			
多発性嚢胞腎	92	19	1			49	15	10	1	187	1	84	272
(%)	(49.2)	(10.2)	(0.5)			(26.2)	(8.0)	(5.3)	(0.5)	(100.0)			
腎硬化症	311	75	20		16	205	74	63		764	2	340	1,106
(%)	(40.7)	(9.8)	(2.6)		(2.1)	(26.8)	(9.7)	(8.2)		(100.0)			
悪性高血圧	32	7	1		1	17	8	7		73	1	32	106
(%)	(43.8)	(9.6)	(1.4)		(1.4)	(23.3)	(11.0)	(9.6)		(100.0)			
糖尿病性腎症	591	182	54		22	547	234	215		1,845	5	896	2,746
(%)	(32.0)	(9.9)	(2.9)		(1.2)	(29.6)	(12.7)	(11.7)		(100.0)			
SLE腎炎	11	3	4			7	8	2		35		20	55
(%)	(31.4)	(8.6)	(11.4)			(20.0)	(22.9)	(5.7)		(100.0)			
アミロイド腎	4		1			4	2			11		5	16
(%)	(36.4)		(9.1)			(36.4)	(18.2)			(100.0)			
痛風腎	8	2				1	2	1		14		9	23
(%)	(57.1)	(14.3)				(7.1)	(14.3)	(7.1)		(100.0)			
先天性代謝異常による腎不全	3	3	1			5	2			14		3	17
(%)	(21.4)	(21.4)	(7.1)			(35.7)	(14.3)			(100.0)			
腎・尿路結核												1	1
(%)													
腎・尿路結石	7	1				3	1	4		16		5	21
(%)	(43.8)	(6.3)				(18.8)	(6.3)	(25.0)		(100.0)			
腎・尿路腫瘍	4		2			2	1			9		3	12
(%)	(44.4)		(22.2)			(22.2)	(11.1)			(100.0)			
閉塞性尿路障害	8	2				2	1	2		15		5	20
(%)	(53.3)	(13.3)				(13.3)	(6.7)	(13.3)		(100.0)			
骨髄腫	1		1				1	3		6		1	7
(%)	(16.7)		(16.7)				(16.7)	(50.0)		(100.0)			
腎形成不全	17	7	5	1		5	2	3		40		25	65
(%)	(42.5)	(17.5)	(12.5)	(2.5)		(12.5)	(5.0)	(7.5)		(100.0)			
不明	256	63	15	1	4	140	66	39		584	1	304	889
(%)	(43.8)	(10.8)	(2.6)	(0.2)	(0.7)	(24.0)	(11.3)	(6.7)		(100.0)			
再導入	15	6	1			11	6	6		45		20	65
(%)	(33.3)	(13.3)	(2.2)			(24.4)	(13.3)	(13.3)		(100.0)			
その他	59	29	2	1	3	48	11	13	5	171		100	271
(%)	(34.5)	(17.0)	(1.2)	(0.6)	(1.8)	(28.1)	(6.4)	(7.6)	(2.9)	(100.0)			
合計	2,186	674	161	5	59	1,549	688	527	7	5,856	16	2,743	8,615
(%)	(37.3)	(11.5)	(2.7)	(0.1)	(1.0)	(26.5)	(11.7)	(9.0)	(0.1)	(100.0)			
記載なし													
(%)													
総計	2,186	674	161	5	59	1,549	688	527	7	5,856	16	2,743	8,615
(%)	(37.3)	(11.5)	(2.7)	(0.1)	(1.0)	(26.5)	(11.7)	(9.0)	(0.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D5215##

(C)Japanese Society for Dialysis Therapy

2016年調査