

表 1461 APD使用の有無 原疾患別（腹膜透析患者）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,032	765	1,797	4	865	2,666
(%)	(57.4)	(42.6)	(100.0)			
慢性腎盂腎炎	40	35	75	1	21	97
(%)	(53.3)	(46.7)	(100.0)			
急速進行性糸球体腎炎	21	20	41	1	19	61
(%)	(51.2)	(48.8)	(100.0)			
妊娠腎/妊娠中毒症	13	7	20		4	24
(%)	(65.0)	(35.0)	(100.0)			
その他の分類不能の腎炎	21	27	48	1	26	75
(%)	(43.8)	(56.3)	(100.0)			
多発性嚢胞腎	89	99	188		84	272
(%)	(47.3)	(52.7)	(100.0)			
腎硬化症	434	309	743	1	362	1,106
(%)	(58.4)	(41.6)	(100.0)			
悪性高血圧	40	30	70		36	106
(%)	(57.1)	(42.9)	(100.0)			
糖尿病性腎症	1,021	790	1,811	2	933	2,746
(%)	(56.4)	(43.6)	(100.0)			
SLE腎炎	20	15	35		20	55
(%)	(57.1)	(42.9)	(100.0)			
アミロイド腎	6	5	11		5	16
(%)	(54.5)	(45.5)	(100.0)			
痛風腎	8	7	15		8	23
(%)	(53.3)	(46.7)	(100.0)			
先天性代謝異常による腎不全	5	9	14		3	17
(%)	(35.7)	(64.3)	(100.0)			
腎・尿路結核					1	1
(%)						
腎・尿路結石	8	6	14		7	21
(%)	(57.1)	(42.9)	(100.0)			
腎・尿路腫瘍	5	4	9		3	12
(%)	(55.6)	(44.4)	(100.0)			
閉塞性尿路障害	9	7	16		4	20
(%)	(56.3)	(43.8)	(100.0)			
骨髄腫	2	3	5		2	7
(%)	(40.0)	(60.0)	(100.0)			
腎形成不全	8	33	41		24	65
(%)	(19.5)	(80.5)	(100.0)			
不明	342	219	561	3	325	889
(%)	(61.0)	(39.0)	(100.0)			
再導入	27	20	47		18	65
(%)	(57.4)	(42.6)	(100.0)			
その他	87	82	169		102	271
(%)	(51.5)	(48.5)	(100.0)			
合計	3,238	2,492	5,730	13	2,872	8,615
(%)	(56.5)	(43.5)	(100.0)			
記載なし						
(%)						
総計	3,238	2,492	5,730	13	2,872	8,615
(%)	(56.5)	(43.5)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D5615##

(C)Japanese Society for Dialysis Therapy

2016年調査