

表 1487 PD透析液交換方法 原疾患別（腹膜透析患者）

原疾患	完全手動式	バッグ交換デバイス			合計	不明	記載なし	総計
		紫外線	熱式無菌 接合装置	左記以外、 半手動も 含む				
慢性糸球体腎炎	540	913	267	58	1,778	9	879	2,666
(%)	(30.4)	(51.3)	(15.0)	(3.3)	(100.0)			
慢性腎盂腎炎	18	35	20	3	76		21	97
(%)	(23.7)	(46.1)	(26.3)	(3.9)	(100.0)			
急速進行性糸球体腎炎	10	18	12	2	42		19	61
(%)	(23.8)	(42.9)	(28.6)	(4.8)	(100.0)			
妊娠腎/妊娠中毒症	10	7	2	1	20		4	24
(%)	(50.0)	(35.0)	(10.0)	(5.0)	(100.0)			
その他の分類不能の腎炎	22	20	4	1	47	1	27	75
(%)	(46.8)	(42.6)	(8.5)	(2.1)	(100.0)			
多発性嚢胞腎	56	94	27	8	185		87	272
(%)	(30.3)	(50.8)	(14.6)	(4.3)	(100.0)			
腎硬化症	209	365	137	24	735	4	367	1,106
(%)	(28.4)	(49.7)	(18.6)	(3.3)	(100.0)			
悪性高血圧	20	40	9	1	70		36	106
(%)	(28.6)	(57.1)	(12.9)	(1.4)	(100.0)			
糖尿病性腎症	413	1,055	286	45	1,799	8	939	2,746
(%)	(23.0)	(58.6)	(15.9)	(2.5)	(100.0)			
SLE腎炎	7	22	4		33	1	21	55
(%)	(21.2)	(66.7)	(12.1)		(100.0)			
アミロイド腎	2	8	1		11		5	16
(%)	(18.2)	(72.7)	(9.1)		(100.0)			
痛風腎	10	2	2	1	15		8	23
(%)	(66.7)	(13.3)	(13.3)	(6.7)	(100.0)			
先天性代謝異常による腎不全		11	2	1	14		3	17
(%)		(78.6)	(14.3)	(7.1)	(100.0)			
腎・尿路結核							1	1
(%)								
腎・尿路結石	6	5	3		14		7	21
(%)	(42.9)	(35.7)	(21.4)		(100.0)			
腎・尿路腫瘍	2	5		1	8		4	12
(%)	(25.0)	(62.5)		(12.5)	(100.0)			
閉塞性尿路障害	5	9	2		16		4	20
(%)	(31.3)	(56.3)	(12.5)		(100.0)			
骨髄腫	1	3	1		5		2	7
(%)	(20.0)	(60.0)	(20.0)		(100.0)			
腎形成不全	14	19	5	3	41		24	65
(%)	(34.1)	(46.3)	(12.2)	(7.3)	(100.0)			
不明	177	276	95	12	560	3	326	889
(%)	(31.6)	(49.3)	(17.0)	(2.1)	(100.0)			
再導入	14	20	12	1	47		18	65
(%)	(29.8)	(42.6)	(25.5)	(2.1)	(100.0)			
その他	41	101	22	3	167	1	103	271
(%)	(24.6)	(60.5)	(13.2)	(1.8)	(100.0)			
合計	1,577	3,028	913	165	5,683	27	2,905	8,615
(%)	(27.7)	(53.3)	(16.1)	(2.9)	(100.0)			
記載なし								
(%)								
総計	1,577	3,028	913	165	5,683	27	2,905	8,615
(%)	(27.7)	(53.3)	(16.1)	(2.9)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D5815##

(C)Japanese Society for Dialysis Therapy

2016年調査