

表 1493 PD透析液交換方法 原疾患別(腹膜透析単独患者)

原疾患	完全手動式	バッグ交換デバイス			合計	不明	記載なし	総計
		紫外線	熱式無菌接合装置	左記以外、半自動も含む				
慢性糸球体腎炎	398	707	221	37	1,363	6	641	2,010
(%)	(29.2)	(51.9)	(16.2)	(2.7)	(100.0)			
慢性腎孟腎炎	12	29	16	3	60	21		81
(%)	(20.0)	(48.3)	(26.7)	(5.0)	(100.0)			
急速進行性糸球体腎炎	9	16	10	2	37	16		53
(%)	(24.3)	(43.2)	(27.0)	(5.4)	(100.0)			
妊娠腎/妊娠中毒症	7	5	2		14	3		17
(%)	(50.0)	(35.7)	(14.3)		(100.0)			
その他の分類不能の腎炎	19	14	3		36	1	17	54
(%)	(52.8)	(38.9)	(8.3)		(100.0)			
多発性囊胞腎	47	80	23	6	156	72		228
(%)	(30.1)	(51.3)	(14.7)	(3.8)	(100.0)			
腎硬化症	174	330	114	21	639	2	302	943
(%)	(27.2)	(51.6)	(17.8)	(3.3)	(100.0)			
悪性高血圧	15	33	9		57	30		87
(%)	(26.3)	(57.9)	(15.8)		(100.0)			
糖尿病性腎症	345	860	246	40	1,491	7	749	2,247
(%)	(23.1)	(57.7)	(16.5)	(2.7)	(100.0)			
SLE腎炎	7	18	3		28	18		46
(%)	(25.0)	(64.3)	(10.7)		(100.0)			
アミロイド腎	2	7	1		10	4		14
(%)	(20.0)	(70.0)	(10.0)		(100.0)			
痛風腎	6	2	1	1	10	5		15
(%)	(60.0)	(20.0)	(10.0)	(10.0)	(100.0)			
先天性代謝異常による腎不全		9	2	1	12	3		15
(%)		(75.0)	(16.7)	(8.3)	(100.0)			
腎・尿路結核						1		1
(%)								
腎・尿路結石	4	3	3		10	7		17
(%)	(40.0)	(30.0)	(30.0)		(100.0)			
腎・尿路腫瘍	1	5		1	7	2		9
(%)	(14.3)	(71.4)		(14.3)	(100.0)			
閉塞性尿路障害	4	8	2		14	3		17
(%)	(28.6)	(57.1)	(14.3)		(100.0)			
骨髓腫	1	3	1		5	2		7
(%)	(20.0)	(60.0)	(20.0)		(100.0)			
腎形成不全	11	18	4	3	36	22		58
(%)	(30.6)	(50.0)	(11.1)	(8.3)	(100.0)			
不明	150	232	84	10	476	2	258	736
(%)	(31.5)	(48.7)	(17.6)	(2.1)	(100.0)			
再導入	12	15	10	1	38	12		50
(%)	(31.6)	(39.5)	(26.3)	(2.6)	(100.0)			
その他	34	84	20	3	141	1	84	226
(%)	(24.1)	(59.6)	(14.2)	(2.1)	(100.0)			
合計	1,258	2,478	775	129	4,640	19	2,272	6,931
(%)	(27.1)	(53.4)	(16.7)	(2.8)	(100.0)			
記載なし								
(%)								
総計	1,258	2,478	775	129	4,640	19	2,272	6,931
(%)	(27.1)	(53.4)	(16.7)	(2.8)	(100.0)			

数値下のカッコ内は行方向の合計に対する%です。

##D5824##

(C)Japanese Society for Dialysis Therapy

2016年調査