

表 1534 PET Cr D/P比 原疾患別 (腹膜透析単独患者)

原疾患	0.5未満	0.5~	0.65~	0.81~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	92	309	331	100	832	1,178	2,010	0.65	0.14
(%)	(11.1)	(37.1)	(39.8)	(12.0)	(100.0)				
慢性腎孟腎炎	8	13	15	7	43	38	81	0.63	0.18
(%)	(18.6)	(30.2)	(34.9)	(16.3)	(100.0)				
急速進行性糸球体腎炎	2	7	7	2	18	35	53	0.66	0.14
(%)	(11.1)	(38.9)	(38.9)	(11.1)	(100.0)				
妊娠腎/妊娠中毒症	2	6	2		10	7	17	0.58	0.09
(%)	(20.0)	(60.0)	(20.0)		(100.0)				
その他の分類不能の腎炎	7	7	4	4	22	32	54	0.60	0.20
(%)	(31.8)	(31.8)	(18.2)	(18.2)	(100.0)				
多発性囊胞腎	11	27	40	15	93	135	228	0.66	0.17
(%)	(11.8)	(29.0)	(43.0)	(16.1)	(100.0)				
腎硬化症	30	120	149	60	359	584	943	0.68	0.14
(%)	(8.4)	(33.4)	(41.5)	(16.7)	(100.0)				
悪性高血圧	4	7	19	4	34	53	87	0.68	0.13
(%)	(11.8)	(20.6)	(55.9)	(11.8)	(100.0)				
糖尿病性腎症	46	271	362	134	813	1,434	2,247	0.68	0.14
(%)	(5.7)	(33.3)	(44.5)	(16.5)	(100.0)				
SLE腎炎	1	4	9	1	15	31	46	0.69	0.12
(%)	(6.7)	(26.7)	(60.0)	(6.7)	(100.0)				
アミロイド腎	1		2	2	5	9	14	0.72	0.17
(%)	(20.0)		(40.0)	(40.0)	(100.0)				
痛風腎	2	1	2	2	7	8	15	0.66	0.14
(%)	(28.6)	(14.3)	(28.6)	(28.6)	(100.0)				
先天性代謝異常による腎不全		3	1	1	5	10	15	0.67	0.11
(%)		(60.0)	(20.0)	(20.0)	(100.0)				
腎・尿路結核						1	1		
(%)									
腎・尿路結石		2	3		5	12	17	0.67	0.10
(%)		(40.0)	(60.0)		(100.0)				
腎・尿路腫瘍		1		1	2	7	9	0.68	0.23
(%)		(50.0)		(50.0)	(100.0)				
閉塞性尿路障害		9		1	10	7	17	0.60	0.09
(%)		(90.0)		(10.0)	(100.0)				
骨髄腫		1	1		2	5	7	0.68	0.13
(%)		(50.0)	(50.0)		(100.0)				
腎形成不全	3	11	6	1	21	37	58	0.61	0.11
(%)	(14.3)	(52.4)	(28.6)	(4.8)	(100.0)				
不明	28	80	98	39	245	491	736	0.66	0.14
(%)	(11.4)	(32.7)	(40.0)	(15.9)	(100.0)				
再導入	2	8	10	1	21	29	50	0.63	0.10
(%)	(9.5)	(38.1)	(47.6)	(4.8)	(100.0)				
その他	13	36	31	8	88	138	226	0.63	0.13
(%)	(14.8)	(40.9)	(35.2)	(9.1)	(100.0)				
合計	252	923	1,092	383	2,650	4,281	6,931	0.66	0.14
(%)	(9.5)	(34.8)	(41.2)	(14.5)	(100.0)				
記載なし									
(%)									
総計	252	923	1,092	383	2,650	4,281	6,931	0.66	0.14
(%)	(9.5)	(34.8)	(41.2)	(14.5)	(100.0)				

数値下の括弧内は行方向の合計に対する%です。

##D6125##

(C)Japanese Society for Dialysis Therapy

2016年調査