

表 1550 残腎Kt/V 原疾患別（腹膜透析単独患者）

| 原疾患           | 0.1未満  | 0.1～   | 0.4～   | 0.8～   | 1.2～   | 1.7～   | 2.0～  | 2.4～   | 合計      | 記載なし  | 総計    | 平均   | 標準偏差 |
|---------------|--------|--------|--------|--------|--------|--------|-------|--------|---------|-------|-------|------|------|
| 慢性糸球体腎炎       | 114    | 155    | 174    | 115    | 65     | 12     | 11    | 5      | 651     | 1,359 | 2,010 | 0.59 | 0.57 |
| (%)           | (17.5) | (23.8) | (26.7) | (17.7) | (10.0) | (1.8)  | (1.7) | (0.8)  | (100.0) |       |       |      |      |
| 慢性腎盂腎炎        | 4      | 5      | 8      | 7      | 3      | 2      | 1     |        | 30      | 51    | 81    | 0.70 | 0.58 |
| (%)           | (13.3) | (16.7) | (26.7) | (23.3) | (10.0) | (6.7)  | (3.3) |        | (100.0) |       |       |      |      |
| 急速進行性糸球体腎炎    | 6      | 6      | 4      | 1      |        |        | 1     |        | 18      | 35    | 53    | 0.33 | 0.49 |
| (%)           | (33.3) | (33.3) | (22.2) | (5.6)  |        |        | (5.6) |        | (100.0) |       |       |      |      |
| 妊娠腎/妊娠中毒症     | 1      | 3      |        | 1      | 1      |        |       |        | 6       | 11    | 17    | 0.52 | 0.60 |
| (%)           | (16.7) | (50.0) |        | (16.7) | (16.7) |        |       |        | (100.0) |       |       |      |      |
| その他の分類不能の腎炎   | 1      | 4      | 10     | 3      | 1      |        |       |        | 19      | 35    | 54    | 0.52 | 0.35 |
| (%)           | (5.3)  | (21.1) | (52.6) | (15.8) | (5.3)  |        |       |        | (100.0) |       |       |      |      |
| 多発性嚢胞腎        | 8      | 19     | 24     | 15     | 11     |        |       |        | 77      | 151   | 228   | 0.61 | 0.44 |
| (%)           | (10.4) | (24.7) | (31.2) | (19.5) | (14.3) |        |       |        | (100.0) |       |       |      |      |
| 腎硬化症          | 31     | 57     | 88     | 55     | 27     | 8      | 6     | 3      | 275     | 668   | 943   | 0.66 | 0.54 |
| (%)           | (11.3) | (20.7) | (32.0) | (20.0) | (9.8)  | (2.9)  | (2.2) | (1.1)  | (100.0) |       |       |      |      |
| 悪性高血圧         | 6      | 5      | 3      | 3      | 3      |        |       |        | 20      | 67    | 87    | 0.47 | 0.49 |
| (%)           | (30.0) | (25.0) | (15.0) | (15.0) | (15.0) |        |       |        | (100.0) |       |       |      |      |
| 糖尿病性腎症        | 80     | 145    | 182    | 120    | 87     | 12     | 11    | 8      | 645     | 1,602 | 2,247 | 0.67 | 0.61 |
| (%)           | (12.4) | (22.5) | (28.2) | (18.6) | (13.5) | (1.9)  | (1.7) | (1.2)  | (100.0) |       |       |      |      |
| SLE腎炎         | 3      | 5      | 4      | 2      |        |        |       |        | 14      | 32    | 46    | 0.40 | 0.34 |
| (%)           | (21.4) | (35.7) | (28.6) | (14.3) |        |        |       |        | (100.0) |       |       |      |      |
| アミロイド腎        | 1      | 1      | 2      |        |        |        |       |        | 4       | 10    | 14    | 0.25 | 0.24 |
| (%)           | (25.0) | (25.0) | (50.0) |        |        |        |       |        | (100.0) |       |       |      |      |
| 痛風腎           | 1      | 1      | 1      | 1      | 1      |        |       |        | 5       | 10    | 15    | 0.56 | 0.51 |
| (%)           | (20.0) | (20.0) | (20.0) | (20.0) | (20.0) |        |       |        | (100.0) |       |       |      |      |
| 先天性代謝異常による腎不全 |        |        | 1      | 2      | 1      |        |       |        | 4       | 11    | 15    | 0.90 | 0.35 |
| (%)           |        |        | (25.0) | (50.0) | (25.0) |        |       |        | (100.0) |       |       |      |      |
| 腎・尿路結核        |        |        |        |        |        |        |       |        |         | 1     | 1     |      |      |
| (%)           |        |        |        |        |        |        |       |        |         |       |       |      |      |
| 腎・尿路結石        | 1      | 2      | 2      |        |        |        |       |        | 5       | 12    | 17    | 0.26 | 0.23 |
| (%)           | (20.0) | (40.0) | (40.0) |        |        |        |       |        | (100.0) |       |       |      |      |
| 腎・尿路腫瘍        | 1      |        |        | 1      |        |        |       |        | 2       | 7     | 9     | 0.55 | 0.78 |
| (%)           | (50.0) |        |        | (50.0) |        |        |       |        | (100.0) |       |       |      |      |
| 閉塞性尿路障害       | 1      |        |        | 2      | 2      | 2      |       | 1      | 8       | 9     | 17    | 1.44 | 0.87 |
| (%)           | (12.5) |        |        | (25.0) | (25.0) | (25.0) |       | (12.5) | (100.0) |       |       |      |      |
| 骨髄腫           |        | 1      |        |        | 1      |        |       |        | 2       | 5     | 7     | 0.75 | 0.92 |
| (%)           |        | (50.0) |        |        | (50.0) |        |       |        | (100.0) |       |       |      |      |
| 腎形成不全         | 4      | 2      | 4      |        | 3      | 1      | 1     | 1      | 16      | 42    | 58    | 0.85 | 0.86 |
| (%)           | (25.0) | (12.5) | (25.0) |        | (18.8) | (6.3)  | (6.3) | (6.3)  | (100.0) |       |       |      |      |
| 不明            | 27     | 42     | 48     | 37     | 15     | 6      | 1     | 3      | 179     | 557   | 736   | 0.65 | 0.71 |
| (%)           | (15.1) | (23.5) | (26.8) | (20.7) | (8.4)  | (3.4)  | (0.6) | (1.7)  | (100.0) |       |       |      |      |
| 再導入           | 4      | 6      | 2      | 1      | 3      | 1      | 1     |        | 18      | 32    | 50    | 0.62 | 0.69 |
| (%)           | (22.2) | (33.3) | (11.1) | (5.6)  | (16.7) | (5.6)  | (5.6) |        | (100.0) |       |       |      |      |
| その他           | 13     | 15     | 17     | 16     | 5      | 1      | 2     | 3      | 72      | 154   | 226   | 0.69 | 0.69 |
| (%)           | (18.1) | (20.8) | (23.6) | (22.2) | (6.9)  | (1.4)  | (2.8) | (4.2)  | (100.0) |       |       |      |      |
| 合計            | 307    | 474    | 574    | 382    | 229    | 45     | 35    | 24     | 2,070   | 4,861 | 6,931 | 0.64 | 0.60 |
| (%)           | (14.8) | (22.9) | (27.7) | (18.5) | (11.1) | (2.2)  | (1.7) | (1.2)  | (100.0) |       |       |      |      |
| 記載なし          |        |        |        |        |        |        |       |        |         |       |       |      |      |
| (%)           |        |        |        |        |        |        |       |        |         |       |       |      |      |
| 総計            | 307    | 474    | 574    | 382    | 229    | 45     | 35    | 24     | 2,070   | 4,861 | 6,931 | 0.64 | 0.60 |
| (%)           | (14.8) | (22.9) | (27.7) | (18.5) | (11.1) | (2.2)  | (1.7) | (1.2)  | (100.0) |       |       |      |      |

数値下のかっこ内は行方向の合計に対する％です。  
##D6225##  
(C)Japanese Society for Dialysis Therapy  
2016年調査