

表 1583 腹膜炎罹患回数(回/年) 年齢(歳)別 (腹膜透析単独患者)

年齢	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
15歳未満	50	8				1					59		26	85
(%)	(84.7)	(13.6)				(1.7)					(100.0)			
15歳～	47	3		2							52		27	79
(%)	(90.4)	(5.8)		(3.8)							(100.0)			
30歳～	264	21	3	1							289		170	459
(%)	(91.3)	(7.3)	(1.0)	(0.3)							(100.0)			
45歳～	920	124	20	2	1		1		1		1,069		581	1,650
(%)	(86.1)	(11.6)	(1.9)	(0.2)	(0.1)		(0.1)		(0.1)		(100.0)			
60歳～	1,710	215	48	14	2	1					1,990		1,006	2,996
(%)	(85.9)	(10.8)	(2.4)	(0.7)	(0.1)	(0.1)					(100.0)			
75歳～	820	150	24	4	1						999		543	1,542
(%)	(82.1)	(15.0)	(2.4)	(0.4)	(0.1)						(100.0)			
90歳～	54	9	4								67		53	120
(%)	(80.6)	(13.4)	(6.0)								(100.0)			
合計	3,865	530	99	23	4	2	1		1		4,525		2,406	6,931
(%)	(85.4)	(11.7)	(2.2)	(0.5)	(0.1)	(0.0)	(0.0)		(0.0)		(100.0)			
不明														
(%)														
記載なし														
(%)														
総計	3,865	530	99	23	4	2	1		1		4,525		2,406	6,931
(%)	(85.4)	(11.7)	(2.2)	(0.5)	(0.1)	(0.0)	(0.0)		(0.0)		(100.0)			
平均	63.63	65.52	67.94	61.87	65.25	39.50	53.00		54.00		63.92		64.00	63.95
標準偏差	14.95	14.87	11.77	17.54	14.27	44.55					14.93		15.08	14.98

数値下のかっこ内は行方向の合計に対する%です。

##D6521##

(C)Japanese Society for Dialysis Therapy

2016年調査