

表 1594 出口部感染罹患回数(回/年) 透析歴(年)別 (腹膜透析患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,930	321	82	31	10	2	1			1	2,378	1	1,239	3,618
(%)	(81.2)	(13.5)	(3.4)	(1.3)	(0.4)	(0.1)	(0.0)			(0.0)	(100.0)			
2年～	1,499	295	83	35	10	3	4	2			1,931	1	1,052	2,984
(%)	(77.6)	(15.3)	(4.3)	(1.8)	(0.5)	(0.2)	(0.2)	(0.1)			(100.0)			
5年～	791	110	38	13		1	1				954		621	1,575
(%)	(82.9)	(11.5)	(4.0)	(1.4)		(0.1)	(0.1)				(100.0)			
10年～	151	20	4	2	1						178	1	110	289
(%)	(84.8)	(11.2)	(2.2)	(1.1)	(0.6)						(100.0)			
15年～	48	3	3	1	1						56		29	85
(%)	(85.7)	(5.4)	(5.4)	(1.8)	(1.8)						(100.0)			
20年～	15	2	2	2							21		7	28
(%)	(71.4)	(9.5)	(9.5)	(9.5)							(100.0)			
25年～	2										2		7	9
(%)	(100.0)										(100.0)			
30年～	3										3		3	6
(%)	(100.0)										(100.0)			
35年～												1		1
(%)														
40年～	2										2			2
(%)	(100.0)										(100.0)			
合計	4,441	751	212	84	22	4	7	3		1	5,525	3	3,069	8,597
(%)	(80.4)	(13.6)	(3.8)	(1.5)	(0.4)	(0.1)	(0.1)	(0.1)		(0.0)	(100.0)			
不明	3										3		15	18
(%)	(100.0)										(100.0)			
記載なし														
(%)														
総計	4,444	751	212	84	22	4	7	3		1	5,528	3	3,084	8,615
(%)	(80.4)	(13.6)	(3.8)	(1.5)	(0.4)	(0.1)	(0.1)	(0.1)		(0.0)	(100.0)			
平均	2.96	2.70	3.13	3.26	2.91	3.50	2.86	2.00			2.93	5.00	3.19	3.03
標準偏差	3.52	2.91	3.45	3.96	3.75	1.29	1.57	1.00			3.45	5.57	3.63	3.52

数値下のかっこ内は行方向の合計に対する%です。

##D6615##

(C)Japanese Society for Dialysis Therapy

2016年調査