

表 1601 出口部感染罹患回数(回/年) 透析歴(年)別 (腹膜透析単独患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,829	305	80	27	10					1	2,252	1	1,153	3,406
(%)	(81.2)	(13.5)	(3.6)	(1.2)	(0.4)					(0.0)	(100.0)			
2年～	1,215	247	67	25	8	2	2	2			1,568		816	2,384
(%)	(77.5)	(15.8)	(4.3)	(1.6)	(0.5)	(0.1)	(0.1)	(0.1)			(100.0)			
5年～	466	78	18	5			1				568		372	940
(%)	(82.0)	(13.7)	(3.2)	(0.9)			(0.2)				(100.0)			
10年～	74	6	1								81		46	127
(%)	(91.4)	(7.4)	(1.2)								(100.0)			
15年～	22		1	1							24		16	40
(%)	(91.7)		(4.2)	(4.2)							(100.0)			
20年～	7	1	1	1							10		6	16
(%)	(70.0)	(10.0)	(10.0)	(10.0)							(100.0)			
25年～	1										1		3	4
(%)	(100.0)										(100.0)			
30年～	2										2			2
(%)	(100.0)										(100.0)			
35年～													1	1
(%)														
40年～														
(%)														
合計	3,616	637	168	59	18	2	3	2		1	4,506	1	2,413	6,920
(%)	(80.2)	(14.1)	(3.7)	(1.3)	(0.4)	(0.0)	(0.1)	(0.0)		(0.0)	(100.0)			
不明	2										2		9	11
(%)	(100.0)										(100.0)			
記載なし														
(%)														
総計	3,618	637	168	59	18	2	3	2		1	4,508	1	2,422	6,931
(%)	(80.3)	(14.1)	(3.7)	(1.3)	(0.4)	(0.0)	(0.1)	(0.0)		(0.0)	(100.0)			
平均	2.34	2.23	2.36	2.44	1.72	3.00	4.00	2.50			2.32		2.58	2.41
標準偏差	2.92	2.34	2.68	3.41	1.36	1.41	1.00	0.71			2.84		3.11	2.94

数値下のかっこ内は行方向の合計に対する%です。

##D6624##

(C)Japanese Society for Dialysis Therapy

2016年調査