

文 献

ケース 1

- 1) 児玉和久, 坂口明, 大和眞史, ほか: 頸動脈洞症候群の1例. その発症メカニズムについての考察. 心臓 25 (Suppl 3): 46-49, 1993.
- 2) Luck JC, Hoover RJ, Biederman RW, et al.: Observation on carotid sinus hypersensitivity from direct intraneural recordings of sympathetic nerve traffic. *Am J Cardiol* 77: 1362-1365, 1996.
- 3) Mano T: Sympathetic nerve mechanisms of human adaptation to environment - Findings obtained by recent microneurographic studies. *Environ Med* 34: 1-35, 1990.
- 4) Matsukawa T, Sugiyama Y, Watanabe T, et al.: Gender difference in age-related changes in muscle sympathetic nerve activity in healthy subjects. *Am J Physiol Regul Integr Comp Physiol* 275: R1600-R1604, 1998.
- 5) McIntosh S, Costa DD, Kenny RA: Outcome of an integrated approach to the investigation of dizziness, falls and syncope in elderly patients referred to a syncope clinic. *Age Ageing* 22: 53-58, 1993.
- 6) O'Mahony D: Pathophysiology of carotid sinus hypersensitivity in elderly patients. *Lancet* 346: 950-952, 1995.
- 7) Sugiyama Y, Matsukawa T, Shamsuzzaman ASM, et al.: Delayed and diminished pressor response to muscle sympathetic nerve activity in the elderly. *J Appl Physiol* 80: 869-875, 1996.
- 8) Yatomi A, Iguchi A, Uemura K, et al.: A rare case of recurrent vasodepressive attacks of 2-hour duration: analysis of the mechanism by muscle sympathetic nerve activity recording. *Clin Cardiol* 12: 164-168, 1989.

ケース 2

- 1) Duckworth WC, Jallepalli P, Solomon SS, et al.: Glucose intolerance in spinal cord injury. *Arch Phys Med Rehabil* 64: 107-110, 1983.
- 2) Duckworth WC, Solomon SS, Jallepalli P: Glucose intolerance due to insulin resistance in patients with spinal cord injuries. *Diabetes* 29: 906-910, 1980.
- 3) 平山正昭, 古池保雄, 金桶吉起, ほか: 自律神経機能不全症における食事性低血圧発現の病態 (3) - β_1 刺激薬と α_1 刺激薬併用による治療効果 - . 自律神経 29: 383-388, 1992.
- 4) Mathias C, da Costa D, Bannister R: Postcibal hypotension in autonomic disorders. Bannister R (ed), *Autonomic failure. A textbook of clinical disorders of the autonomic nervous system*, 2nd ed., p.367-380, Oxford University Press, 1988.
- 5) Mathias CJ, Frankel HL, Turner RC, et al.: Physiological responses to insulin hypoglycaemia in spinal man. *Paraplegia* 17: 319-326, 1979.
- 6) 小田嶋奈津, 市川 忠, 古川哲雄, ほか: 高位脊髄損傷患者における経口ぶどう糖負荷試験による血圧の変動. 自律神経 28: 266-269, 1991.

ケース 3

- 1) 長谷川康博, 古池保雄, 松岡幸彦, ほか: Shy-Drager 症候群にみられた“食事性低血圧・失神”とその L-DOPS による治療. 日内会誌 78: 939-943, 1989.

ケース 4

- 1) 長谷川康博, 岡本 進: 著明な食事性低血圧による食後失神を主訴としたパーキンソン病. 臨床神経 32: 1017-1021, 1992.

ケース 5

- 1) Koike Y, Takahashi A: Autonomic dysfunction in Parkinson's disease. *Eur Neurol* 38 (Suppl 2): 8-12, 1997.
- 2) Linden D, Diehl RR, Berlit P: Sympathetic cardiovascular dysfunction in long-standing idiopathic Parkinson's disease. *Clin Auton Res* 7: 311-314, 1997.
- 3) Niimi Y, Ieda T, Hirayama M, et al.: Clinical and physiological characteristics of autonomic failure with Parkinson's disease. *Clin Auton Res* 9: 139-144, 1999.